

Product Sheet

H_CD24 MC38 Cell Line

Catalog number: GM-C15769

Version 3.3.1.260126

Description	H_CD24 MC38 Cell Line is a clonal stable MC38 cell line that constitutively expresses the human CD24 gene, constructed using lentiviral technology.
Quantity	5E6 Cells per vial, 1 mL
Product Format	1 vial of frozen cells
Shipping	Shipped on dry ice
Storage Conditions	Liquid nitrogen immediately upon receipt
Target	Human_CD24
Gene ID/Uniprot ID	P25063-1
Host Cell	MC38
Recovery Medium	DMEM+10% FBS+1% P.S
Growth medium	DMEM+10% FBS+1% P.S+2.5 µg/mL Puromycin
Note	None
Freezing Medium	90% FBS+10% DMSO
Growth properties	Adherent
Growth Conditions	37°C, 5% CO ₂
Mycoplasma Testing	The cell line has been screened to confirm the absence of Mycoplasma species.
Safety considerations	Biosafety Level 2
Note	It is recommended to expand the cell culture and store a minimum of 10 vials at an early passage for potential future use.

Materials

Reagent	Manufacturer/Catalogue No.
DMEM	VivaCell/C3110-0500
Fetal Bovine Serum	ExCell/FSP500
Pen/Strep	Thermo/15140-122
Puromycin	Genomeditech/GM-040401
Anti-CD24 hIgG1 Antibody(LM-109)	Genomeditech/GM-82520AB

Figures




Figure 1 | H_CD24 MC38 Cell Line (Cat. GM-C15769) was determined by flow cytometry using Anti-CD24 hIgG1 Antibody(LM-109) (Cat. GM-82520AB).




Figure 2 | H_CD24 MC38 Cell Line (Cat. GM-C15769) was determined by flow cytometry using Anti-CD24 hIgG1 Antibody(LM-109) (Cat. GM-82520AB).




Figure 3 | Tumor growth curves of H_CD24 MC38 in C57BL/6 mice. H_CD24 MC38 cells (1×10^6 per mouse) were subcutaneously inoculated into C57BL/6 mice (female, 8 weeks old, $n = 3$). Tumor volume was measured twice per week and is presented as mean \pm SEM.




Figure 4 | Body weight changes after implantation of H_CD24 MC38 in C57BL/6 mice. Under the same conditions, body weight was measured twice per week and is presented as mean \pm SEM.

Cell Recovery

Recovery Medium: DMEM+10% FBS+1% P.S

To insure the highest level of viability, thaw the vial and initiate the culture as soon as possible upon receipt. If upon arrival, continued storage of the frozen culture is necessary, it should be stored in liquid nitrogen vapor phase and not at -70°C. Storage at -70°C will result in loss of viability.

- a) Thaw the vial by gentle agitation in a 37°C water bath. To reduce the possibility of contamination, keep the O-ring and cap out of the water. Thawing should be rapid (approximately 2 - 3 minutes).
- b) Remove the vial from the water bath as soon as the contents are thawed, and decontaminate by dipping in or spraying with 70% ethanol. All of the operations from this point on should be carried out under strict aseptic conditions.
- c) Transfer the vial contents to a centrifuge tube containing 5.0 mL complete culture medium and spin at approximately 176 x g for 5 minutes. Discard supernatant.
- d) Resuspend cell pellet with the recommended recovery medium. And dispense into appropriate culture dishes.
- e) Incubate the culture at 37°C in a suitable incubator. A 5% CO₂ in air atmosphere is recommended if using the medium described on this product sheet.

Cell Freezing

Freezing Medium: 90% FBS+10% DMSO

- a) Centrifuge at 176 x g for 3 minutes to collect cells.
- b) Resuspend the cells in pre-cooled freezing medium and adjust the cell density to 5E6 cells/mL.
- c) Aliquot 1 mL into each vial.
- d) Place the vial in a controlled-rate freezing container and store at -80°C for at least 1 day, then transfer to liquid nitrogen as soon as possible.

Cell passage

Growth medium: DMEM+10% FBS+1% P.S+2.5 µg/mL Puromycin

For the first 1 to 2 passages post-resuscitation, use the recovery medium. Once the cells have stabilized, switch to a growth medium.

- a) Remove and discard culture medium.
- b) Briefly rinse the cell layer with PBS to remove all traces of serum that contains trypsin inhibitor.
- c) Add 1.0 mL of 0.25% (w/v) Trypsin-EDTA solution to dish and observe cells under an inverted microscope until cell layer is dispersed (usually within 30 to 60 seconds at 37°C).
- d) Note: To avoid clumping do not agitate the cells by hitting or shaking the flask while waiting for the cells to detach. Cells that are difficult to detach may be placed at 37°C to facilitate dispersal.
- e) Add 2.0 mL of growth medium to mix well and aspirate cells by gently pipetting.
- f) After centrifugation, resuspend the pellet and add appropriate aliquots of the cell suspension to new culture vessels.
- g) Incubate cultures at 37°C.

Subcultivation Ratio: A subcultivation ratio of 1:4 - 1:5 is recommended

Medium Renewal: Every 2 to 3 days

Notes

- a) After the stabilization of the cell condition, there will be fewer dead cells post-passage, the cell growth rate will tend to stabilize, cell morphology will become uniform, and the cells will appear robust.

Sequence

CD24 P25063-1

MGRAMVARGLGLLLLALLLPTQIYSSETTGTSSNNSQSTSNSGLAPNPTNATTKAAGGALQSTASLFVVSL
SLLHLYS*

Related Products

CD24:Siglec10	
Flag-Cynomolgus_CD24 CHO-K1 Cell Line	H_CD24 CHO-K1 Cell Line
H_CD24 HEK-293 Cell Line	H_Siglec10 CHO-K1 Cell Line
MCF-7(CD24-Positive) Luciferase Cell Line	
Anti-CD24 hIgG1 Antibody(LM-109)	
SIGLEC15	
Cynomolgus_SIGLEC15 CHO-K1 Cell Line	H_SIGLEC15 CHO-K1 Cell Line
H_SIGLEC15 HEK-293 Cell Line	H_SIGLEC15 MC38 Cell Line
H_SIGLEC15 U2OS Cell Line	Mouse_SIGLEC15 CHO-K1 Cell Line
Anti-Siglec15 mIgG2a Antibody(5G12)	
SIGLEC9	
Cynomolgus_SIGLEC9 CHO-K1 Cell Line	H_SIGLEC9 CHO-K1 Cell Line
H_SIGLEC9 HEK-293 Cell Line	
Anti-siglec9 mIgG1 Antibody(2D4)	
SIGLEC8	
H_SIGLEC8 CHO-K1 Cell Line	H_SIGLEC8 HEK-293 Cell Line
Olive Baboon_SIGLEC8 CHO-K1 Cell Line	
Anti-H_SIGLEC8 hIgG1 Antibody(1H10)	
SIGLEC3(CD33)	
Cynomolgus_CD33(SIGLEC3) CHO-K1 Cell Line	H_CD33(SIGLEC3) CHO-K1 Cell Line
Anti-H_CD33(siglec3) hIgG4 Antibody(Gemtuzumab)	
In Vivo MAb Isotype Controls	
Biotinylated Human IgE Isotype Control; His-Avi Tag (Anti-RSV)	Human IgG1 (K214R) Isotype Control(Anti-HEL)
Human IgG1 (K214R;D356E;L358M) Isotype Control(Anti-HEL)	Human IgG1 (KRLALAGA) Isotype Control(Anti-HEL)
Human IgG1 Isotype Control(Anti-HEL)	Human IgG1 Isotype Control(Anti-RSV)
Human IgG1 Isotype Control(MOPC-21)	Human IgG1(LALA) Isotype Control(Anti-HEL)
Human IgG1(LALAPG) Isotype Control(Anti-HEL)	Human IgG1(LALAPG;N297A) Isotype Control(Anti-HEL)
Human IgG1(N297A) Isotype Control(Anti-HEL)	Human IgG4(S228P) Isotype Control(Anti-HEL)
Mouse IgG1 (D265A) Isotype Control(Anti-HEL)	Mouse IgG1 Isotype Control(Anti-HEL)
Mouse IgG2a Isotype Control(Anti-HEL)	Mouse IgG2a Isotype Control(Anti-RSV)
Mouse IgG2a(D265A) Isotype Control(Anti-HEL)	Mouse IgG2b Isotype Control(Anti-HEL)
Rat IgG1 Isotype Control(Anti-HEL)	

License Agreement:

By purchasing and using this cell line product, the user voluntarily agrees to accept and abide by the following policies:

- This cell line product is restricted to research use only and shall not be used for any commercial purposes.
- This product is strictly prohibited from being used in the diagnosis or treatment of human or animal diseases, and shall not be directly used in experiments involving humans.
- Users and their contractors engaged for their benefit may use this material and its derivatives only within the agreed research scope; modification of the material is not permitted, nor may it be distributed, sold, transferred, or otherwise provided to any other entity (including affiliates).
- If use beyond the above scope is required, prior written permission from Genomeditech (Shanghai) Co.,Ltd. must be obtained. For details, please contact Genomeditech (Shanghai) Co.,Ltd.