

# Product Sheet


## H\_TLR2 Reporter 293 Cell Line

Catalog number: GM-C15493

Version 3.3.1.260226

TLR2 (Toll-like receptor 2) is a widely distributed transmembrane protein belonging to the Toll-like receptor family and acts as an important pattern recognition receptor (PRR) in the innate immune system. TLR2 is mainly expressed on the surface of immune cells such as monocytes, macrophages, and dendritic cells, and is capable of recognizing various pathogen-associated molecular patterns (PAMPs), such as bacterial lipoproteins, peptidoglycan, and lipopolysaccharides. By binding to its ligands, TLR2 can activate downstream signaling pathways, induce the production of inflammatory cytokines, and promote the initiation and regulation of immune responses. Studies have shown that TLR2 plays a key role not only in defending against infections and clearing pathogens, but is also closely associated with a variety of inflammatory diseases, autoimmune disorders, and tumorigenesis.

H\_TLR2 Reporter 293 Cell Line is a clonal stable 293 cell line constructed using lentiviral technology, constitutive expression of the TLR2 gene, along with signal-dependent expression of a luciferase reporter gene. When Pam3CSK4 binds to TLR2, it activates downstream signaling pathways, leading to the expression of luciferase. The luciferase activity measurement indicates the activation level of the signaling pathway and can thus be used to evaluate the in vitro effects of drugs related to TLR2.



## Specifications

<b>Quantity</b>	5E6 Cells per vial, 1 mL
<b>Product Format</b>	1 vial of frozen cells
<b>Shipping</b>	Shipped on dry ice
<b>Storage Conditions</b>	Liquid nitrogen immediately upon receipt
<b>Recovery Medium</b>	EMEM+10% FBS+1% P.S
<b>Growth medium</b>	EMEM+10% FBS+1% P.S+3 µg/mL Blasticidin+400 µg/mL G418+1.5 µg/mL Puromycin
<b>Note</b>	None
<b>Freezing Medium</b>	90% FBS+10% DMSO
<b>Growth properties</b>	Adherent
<b>Growth Conditions</b>	37°C, 5% CO <sub>2</sub>
<b>Mycoplasma Testing</b>	The cell line has been screened to confirm the absence of Mycoplasma species.
<b>Safety considerations</b>	Biosafety Level 2
<b>Note</b>	It is recommended to expand the cell culture and store a minimum of 10 vials at an early passage for potential future use.

## Materials

Reagent	Manufacturer/Catalogue No.
EMEM	ATCC/30-2003
Fetal Bovine Serum	ExCell/FSP500
Pen/Strep	Thermo/15140-122
Blasticidin	Genomeditech/ <a href="#">GM-040404</a>
G418	Genomeditech/ <a href="#">GM-040402</a>
Puromycin	Genomeditech/ <a href="#">GM-040401</a>
Pam3CSK4 TFA	MCE/HY-P1180A
Tomaralimab (anti-TLR2)	Aladdin/Ab176102
GMOne-Step 2.0 Luciferase Reporter Gene Assay Kit	Genomeditech/ <a href="#">GM-040513</a>

## Figures




Figure 1 | Response to Pam3CSK4(TFA). The H\_TLR2 Reporter 293 Cell Line (Cat. GM-C15493) at a concentration of 2E4 cells/well (96-well format) was stimulated with serial dilutions of Pam3CSK4(TFA) (MCE/HY-P1180A) in assay buffer (EMEM+1% FBS+1% P.S) for 6 hours. The firefly luciferase activity was measured using the Luciferase Reporter Assay Kit (Genomeditech). The maximum induction fold was approximately [119.5]. Data are shown by drug mass concentration.




Figure 2 | Response to Pam3CSK4(TFA). The H\_TLR2 Reporter 293 Cell Line (Cat. GM-C15493) at a concentration of 2E4 cells/well (96-well format) was stimulated with serial dilutions of Pam3CSK4(TFA) (MCE/HY-P1180A) in assay buffer (EMEM+1% FBS+1% P.S) for 6 hours. The firefly luciferase activity was measured using the Luciferase Reporter Assay Kit (Genomeditech). The maximum induction fold was approximately [119.5]. Data are shown by drug molar concentration.




Figure 3 | H\_TLR2 Reporter 293 Cell Line (Cat. GM-C15493) was determined by flow cytometry using Tomaralimab (anti-TLR2) (Aladdin/Ab176102).

## Cell Recovery

Recovery Medium: EMEM+10% FBS+1% P.S

To insure the highest level of viability, thaw the vial and initiate the culture as soon as possible upon receipt. If upon arrival, continued storage of the frozen culture is necessary, it should be stored in liquid nitrogen vapor phase and not at  $-70^{\circ}\text{C}$ . Storage at  $-70^{\circ}\text{C}$  will result in loss of viability.

- Thaw the vial by gentle agitation in a  $37^{\circ}\text{C}$  water bath. To reduce the possibility of contamination, keep the O-ring and cap out of the water. Thawing should be rapid (approximately 2 - 3 minutes).
- Remove the vial from the water bath as soon as the contents are thawed, and decontaminate by dipping in or spraying with 70% ethanol. All of the operations from this point on should be carried out under strict aseptic conditions.
- Transfer the vial contents to a centrifuge tube containing 5.0 mL complete culture medium and spin at approximately  $176 \times g$  for 5 minutes. Discard supernatant.
- Resuspend cell pellet with the recommended recovery medium. And dispense into appropriate culture dishes.
- Incubate the culture at  $37^{\circ}\text{C}$  in a suitable incubator. A 5%  $\text{CO}_2$  in air atmosphere is recommended if using the medium described on this product sheet.

## Cell Freezing

Freezing Medium: 90% FBS+10% DMSO

- Centrifuge at  $176 \times g$  for 3 minutes to collect cells.
- Resuspend the cells in pre-cooled freezing medium and adjust the cell density to  $5 \times 10^6$  cells/mL.
- Aliquot 1 mL into each vial.
- Place the vial in a controlled-rate freezing container and store at  $-80^{\circ}\text{C}$  for at least 1 day, then transfer to liquid nitrogen as soon as possible.

## Cell passage

Growth medium: EMEM+10% FBS+1% P.S+3  $\mu\text{g}/\text{mL}$  Blasticidin+400  $\mu\text{g}/\text{mL}$  G418+1.5  $\mu\text{g}/\text{mL}$  Puromycin

For the first 1 to 2 passages post-resuscitation, use the recovery medium. Once the cells have stabilized, switch to a growth medium.

- Remove and discard culture medium.
- Briefly rinse the cell layer with PBS to remove all traces of serum that contains trypsin inhibitor.
- Add 1.0 mL of 0.25% (w/v) Trypsin-EDTA solution to dish and observe cells under an inverted microscope until cell layer is dispersed (usually within 30 to 60 seconds at 37°C).
- Note: To avoid clumping do not agitate the cells by hitting or shaking the flask while waiting for the cells to detach. Cells that are difficult to detach may be placed at 37°C to facilitate dispersal.
- Add 2.0 mL of growth medium to mix well and aspirate cells by gently pipetting.
- After centrifugation, resuspend the pellet and add appropriate aliquots of the cell suspension to new culture vessels.
- Incubate cultures at 37°C.

**Subcultivation Ratio: A subcultivation ratio of 1:3 - 1:4 is recommended**

**Medium Renewal: Every 2 to 3 days**

## Notes

- Upon initial revival, a higher number of dead cells and poor adherence are observed, which is normal. Adherence typically recovers within 2 - 3 days. After 2 - 3 passages, the proportion of adherent cells increases, and the cells begin to spread normally.
- After each passage, there may be 5 - 10% dead cells; however, as the number of passages increases, the recovery rate accelerates, the proportion of dead cells decreases, and the cell growth rate stabilizes.
- It is recommended to retain cell images after revival and during each observation to assist in assessing cell status. In case of abnormalities, promptly communicate with Genomeditech sales.

## Related Products

TLR7	
<a href="#">H_TLR7 Reporter 293 Cell Line</a>	<a href="#">Mouse_TLR7 Reporter 293 Cell Line</a>
TLR9	
<a href="#">H_TLR9 Reporter 293 Cell Line</a>	<a href="#">Mouse_TLR9 Reporter 293 Cell Line</a>
TLR8	
<a href="#">H_TLR8 Reporter 293 Cell Line</a>	<a href="#">H_TLR8 HEK-293 Cell Line</a>
STING	
<a href="#">H_STING KO THP1 Cell Line</a>	<a href="#">H_STING KO U937 Cell Line</a>
<a href="#">STING KO Reporter THP1 Cell Line</a>	<a href="#">STING Reporter HEK-293 Cell Line</a>
<a href="#">STING Reporter THP1 Cell Line</a>	<a href="#">STING Reporter U937 Cell Line</a>

## License Agreement:

**By purchasing and using this cell line product, the user voluntarily agrees to accept and abide by the following policies:**

- This cell line product is restricted to research use only and shall not be used for any commercial purposes.
- This product is strictly prohibited from being used in the diagnosis or treatment of human or animal diseases, and shall not be directly used in experiments involving humans.
- Users and their contractors engaged for their benefit may use this material and its derivatives only within the agreed research scope; modification of the material is not permitted, nor may it be distributed, sold, transferred, or otherwise provided to any other entity (including affiliates).
- If use beyond the above scope is required, prior written permission from Genomeditech (Shanghai) Co.,Ltd. must be obtained. For details, please contact Genomeditech (Shanghai) Co.,Ltd.

Genomeditech