

Product Sheet

STING Reporter THP1 Cell Line


Catalog number: GM-C21640

Version 3.3.1.260610

STING (Stimulator of Interferon Genes) is a key intracellular receptor involved in the immune response to viral and bacterial infections. It recognizes cyclic dinucleotides like cGAMP in the cytoplasm, which are produced by pathogens or synthesized by host cells during infection. STING activation enhances the production of interferons and inflammatory factors, boosting antiviral and antitumor responses.

The STING signaling pathway is mediated by its interaction with TBK1 and IRF3. When STING binds to cGAMP, it recruits TBK1, which phosphorylates IRF3, activating and translocating it to the nucleus. There, IRF3 promotes the transcription of interferon genes, initiating antiviral responses. STING can also activate the NF- κ B pathway, enhancing inflammation.

STING Reporter THP1 Cell Line is a clonal stable cell line with signal-dependent expression of a luciferase reporter gene constructed using lentiviral technology, and it endogenously expresses STING gene(R232) and IFNAR gene. When ADU-S100 binds to STING, it activates downstream signaling pathways, leading to the expression of luciferase. H-151 can inhibit this signal transmission. The luciferase activity measurement indicates the activation level of the signaling pathway and can thus be used to evaluate the in vitro effects of drugs related to STING.



Specifications

Quantity	5E6 Cells per vial, 1 mL
Product Format	1 vial of frozen cells
Shipping	Shipped on dry ice
Storage Conditions	Liquid nitrogen immediately upon receipt
Recovery Medium	RPMI 1640(ATCC)+10% FBS+1% P.S+0.05 mM β -Me
Growth medium	RPMI 1640(ATCC)+10% FBS+1% P.S+0.05 mM β -Me+2 μ g/mL Blasticidin
Note	Cells should be cultured using ATCC/30-2001 RPMI 1640 medium or Growth medium from Genomeditech. The serum should be ExCell/FSP500 or sourced from Gibco.
Freezing Medium	90% FBS+10% DMSO
Growth properties	Suspension
Growth Conditions	37°C, 5% CO ₂
Mycoplasma Testing	The cell line has been screened to confirm the absence of Mycoplasma species.
Safety considerations	Biosafety Level 2
Note	It is recommended to expand the cell culture and store a minimum of 10 vials at an early passage for potential future use.

Materials

Reagent	Manufacturer/Catalogue No.
RPMI 1640(ATCC)	ATCC/30-2001
Fetal Bovine Serum	ExCell/FSP500
Pen/Strep	Thermo/15140-122
2-Mercaptoethanol(β -Me)	gibco/21985-023
Blasticidin	Genomeditech/ GM-040404
ADU-S100 disodium salt	MCE/HY-12885A
GMOne-Step 2.0 Luciferase Reporter Gene Assay Kit	Genomeditech/ GM-040513

Figures




Figure 1 | Response to ADU-S100 disodium salt. The STING Reporter THP1 Cell Line (Cat. GM-C21640) at a concentration of 1E5 cells/well (96-well format) was stimulated with serial dilutions of ADU-S100 disodium salt (MCE/HY-12885A) in assay buffer (RPMI 1640(ATCC) + 1% FBS + 1% P.S) for 16 hours. The firefly luciferase activity was measured using the Luciferase Reporter Assay Kit (Genomeditech). The maximum induction fold was approximately [3.7]. Data are shown by drug molar concentration.

Cell Recovery

Recovery Medium: RPMI 1640(ATCC)+10% FBS+1% P.S+0.05 mM β -Me

To insure the highest level of viability, thaw the vial and initiate the culture as soon as possible upon receipt. If upon arrival, continued storage of the frozen culture is necessary, it should be stored in liquid nitrogen vapor phase and not at -70°C . Storage at -70°C will result in loss of viability.

- a) Thaw the vial by gentle agitation in a 37°C water bath. To reduce the possibility of contamination, keep the O-ring and cap out of the water. Thawing should be rapid (approximately 2 - 3 minutes).
- b) Remove the vial from the water bath as soon as the contents are thawed, and decontaminate by dipping in or spraying with 70% ethanol. All of the operations from this point on should be carried out under strict aseptic conditions.
- c) Transfer the vial contents to a centrifuge tube containing 5.0 mL complete culture medium. And spin at approximately $176 \times g$ for 5 minutes. Discard supernatant.
- d) Resuspend cell pellet with the recommended complete medium. And dispense the suspension into an appropriate culture flask and initially place the flask in an upright position after thawing.
- e) Incubate the culture at 37°C in a suitable incubator. A 5% CO_2 in air atmosphere is recommended if using the medium described on this product sheet.

Cell Freezing

Freezing Medium: 90% FBS+10% DMSO

- Centrifuge at 176 x g for 3 minutes to collect cells.
- Resuspend the cells in pre-cooled freezing medium and adjust the cell density to 5E6 cells/mL.
- Aliquot 1 mL into each vial.
- Place the vial in a controlled-rate freezing container and store at -80°C for at least 1 day, then transfer to liquid nitrogen as soon as possible.

Cell passage

Growth medium: RPMI 1640(ATCC)+10% FBS+1% P.S+0.05 mM β -Me+2 μ g/mL Blasticidin

During the first two passages after cell thawing, use the recovery medium. Once the cell status stabilizes, switch to growth medium containing antibiotics.

- When the cell density reaches 8E5 cells/mL, subculture the cells. Do not allow the cell density to exceed 1E6 cells/mL.
- It is recommended to use T-25 flasks for subculturing.
- These cells are suspension cells, and it is recommended to use the "half-medium change" method to maintain optimal cell conditions during passaging.
- During passaging, you can directly add fresh growth medium to the culture flask, gently pipette to resuspend the cells, and then transfer the cell suspension to a new T-25 flask for continued culture.

Subcultivation Ratio: Maintain cultures at a cell concentraion between 2.5E5 and 8E5 viable cells/mL.

Medium Renewal: Every 2 to 3 days

Notes

- After thawing, cell growth is slow, and there will be a significant amount of cellular debris in the background. As the cells recover, the background will gradually become cleaner, with a recovery period estimated at 1 to 1.5 weeks.
- These cells are sensitive to cell density, so please ensure that cell density is maintained within an appropriate range during culture and passaging.
- The culture medium for these cells must be supplemented with β -mercaptoethanol. Failure to add this supplement may negatively affect cell status.
- Cells should be cultured using ATCC/30-2001 RPMI 1640 medium or complete medium purchased from Geomeditech. The serum used should be the same as specified in the manual or Gibco serum.

Related Products

TLR2	
H_TLR2 Reporter 293 Cell Line	
TLR7	
H_TLR7 Reporter 293 Cell Line	Mouse_TLR7 Reporter 293 Cell Line
TLR9	

H_TLR9 Reporter 293 Cell Line	Mouse_TLR9 Reporter 293 Cell Line
TLR8	
H_TLR8 Reporter 293 Cell Line	H_TLR8 HEK-293 Cell Line
STING	
H_STING KO THP1 Cell Line	H_STING KO U937 Cell Line
STING KO Reporter THP1 Cell Line	STING Reporter HEK-293 Cell Line
STING Reporter U937 Cell Line	

License Agreement:

By purchasing and using this cell line product, the user voluntarily agrees to accept and abide by the following policies:

- This cell line product is restricted to research use only and shall not be used for any commercial purposes.
- This product is strictly prohibited from being used in the diagnosis or treatment of human or animal diseases, and shall not be directly used in experiments involving humans.
- Users and their contractors engaged for their benefit may use this material and its derivatives only within the agreed research scope; modification of the material is not permitted, nor may it be distributed, sold, transferred, or otherwise provided to any other entity (including affiliates).
- If use beyond the above scope is required, prior written permission from Genomeditech (Shanghai) Co.,Ltd. must be obtained. For details, please contact Genomeditech (Shanghai) Co.,Ltd.