

Product Sheet

Rabbit_CD40L NIH-3T3 Cell Line

Catalog number: GM-C42360

Version 3.3.1.260618

Description	Rabbit_CD40L NIH-3T3 Cell Line is a clonal stable NIH-3T3 cell line that constitutively expresses the Rabbit CD40L gene, constructed using lentiviral technology.
Quantity	5E6 Cells per vial, 1 mL
Product Format	1 vial of frozen cells
Shipping	Shipped on dry ice
Storage Conditions	Liquid nitrogen immediately upon receipt
Target	Rabbit CD40L
Gene ID/Uniprot ID	G1SKP7
Host Cell	NIH-3T3
Recovery Medium	DMEM+10% FBS+1% P.S
Growth medium	DMEM+10% FBS+1% P.S+1 µg/mL Puromycin
Note	None
Freezing Medium	90% FBS+10% DMSO
Growth properties	Adherent
Growth Conditions	37°C, 5% CO ₂
Mycoplasma Testing	The cell line has been screened to confirm the absence of Mycoplasma species.
Safety considerations	Biosafety Level 2
Note	It is recommended to expand the cell culture and store a minimum of 10 vials at an early passage for potential future use.

Materials

Reagent	Manufacturer/Catalogue No.
DMEM	Gibco/C11995500BT
Fetal Bovine Serum	ExCell/FSP500
Pen/Strep	Thermo/15140-122
Puromycin	Genomeditech/GM-040401
Anti-CD40L hIgG1 Reference Antibody (Frebio)	Genomeditech/GM-87982MAB
Anti-His mIgG2a Antibody	Genomeditech/GM-59493AB

Figures




Figure 1 | Rabbit_CD40L NIH-3T3 Cell Line (Cat. GM-C42360) was determined by flow cytometry using Anti-CD40L hIgG1 Reference Antibody (Frebio) (Cat. GM-87982MAB).




Figure 2 | Rabbit_CD40L NIH-3T3 Cell Line (Cat. GM-C42360) was determined by flow cytometry using Anti-His mIgG2a Antibody (Cat. GM-59493AB).

Cell Recovery

Recovery Medium: DMEM+10% FBS+1% P.S

To insure the highest level of viability, thaw the vial and initiate the culture as soon as possible upon receipt. If upon arrival, continued storage of the frozen culture is necessary, it should be stored in liquid nitrogen vapor phase and not at -70°C . Storage at -70°C will result in loss of viability.

- a) Thaw the vial by gentle agitation in a 37°C water bath. To reduce the possibility of contamination, keep the O-ring and cap out of the water. Thawing should be rapid (approximately 2 - 3 minutes).
- b) Remove the vial from the water bath as soon as the contents are thawed, and decontaminate by dipping in or spraying with 70% ethanol. All of the operations from this point on should be carried out under strict aseptic conditions.
- c) Transfer the vial contents to a centrifuge tube containing 5.0 mL complete culture medium and spin at approximately $176 \times g$ for 5 minutes. Discard supernatant.
- d) Resuspend cell pellet with the recommended recovery medium. And dispense into appropriate culture dishes.
- e) Incubate the culture at 37°C in a suitable incubator. A 5% CO_2 in air atmosphere is recommended if using the medium described on this product sheet.

Cell Freezing

Freezing Medium: 90% FBS+10% DMSO

- a) Centrifuge at $176 \times g$ for 3 minutes to collect cells.
- b) Resuspend the cells in pre-cooled freezing medium and adjust the cell density to 5×10^6 cells/mL.
- c) Aliquot 1 mL into each vial.
- d) Place the vial in a controlled-rate freezing container and store at -80°C for at least 1 day, then transfer to liquid nitrogen as soon as possible.

Cell passage

Growth medium: DMEM+10% FBS+1% P.S+1 $\mu\text{g/mL}$ Puromycin

For the first 1 to 2 passages post-resuscitation, use the recovery medium. Once the cells have stabilized, switch to a growth medium.

- a) Subculturing is necessary when the cell density reaches 60%. It is recommended to perform subculturing at a ratio of 1:4 to 1:5 every 2-3 days.
 - b) Remove and discard culture medium.
 - c) Briefly rinse the cell layer with PBS to remove all traces of serum that contains trypsin inhibitor.
 - d) Add 1.0 mL of 0.25% (w/v) Trypsin-EDTA solution to dish and observe cells under an inverted microscope until cell layer is dispersed (usually within 30 seconds at 37°C).
 - e) Note: To avoid clumping do not agitate the cells by hitting or shaking the flask while waiting for the cells to detach. Cells that are difficult to detach may be placed at 37°C to facilitate dispersal.
 - f) Add 2.0 mL of growth medium to mix well and aspirate cells by gently pipetting.
 - g) After centrifugation, resuspend the pellet and add appropriate aliquots of the cell suspension to new culture vessels.
-

h) Incubate cultures at 37°C.

Subcultivation Ratio: A subcultivation ratio of 1:4 - 1:5 is recommended

Medium Renewal: Every 2 to 3 days

Notes

- FBS needs to be heat-inactivated at 56°C for 30 minutes. This process inactivates complement proteins and some viruses, but does not significantly affect the activity of most growth factors and cytokines.
- Cells should never be allowed to become over-confluent. Subculturing should be performed at least twice per week to ensure that cell density does not exceed 80%.

Sequence

CD40L-His [G1SKP7](#)

MIETYSQPTPRSVATGPSVSMKIFMYLLTVFLITQMIGSALFAVYLHRRRLDKIEDERNLHEDFVFMKTIQRCNK
GEGSLSLNCKEIRSQFEGFVKDIMLNKEEPKKEINFEMQKGDQDPQIAAHLISEASSKSSSVLQWAKKGYTT
MSNTLVTLENGKQLKVKRQGFYYIYAQVTFCSNQEPSSQAPFIASLCLKSSGGSERILLRAANARSSSKTCEQ
QSIHLGGVFELQADASVFNVTASQVNHGTGFTSFGLLKLHHHHHH

Related Products

CD40:CD40L	
H_CD40(TNFRSF5) Reporter 293 Cell Line	H_CD40(TNFRSF5) Reporter Jurkat Cell Line
Cynomolgus_CD40 CHO-K1 Cell Line	Cynomolgus_CD40L CHO-K1 Cell Line
H_CD40(TNFRSF5) CHO-K1 Cell Line	H_CD40(TNFRSF5) HEK-293 Cell Line
H_CD40L CHO-K1 Cell Line	H_CD40L HEK-293 Cell Line
Mouse_CD40L CHO-K1 Cell Line	
Anti-CD40 hIgG1 Reference Antibody (Sotibio)	Anti-CD40 hIgG1 Reference Antibody (Tenebio)
Anti-CD40L hIgG1 Reference Antibody (Frebio)	Anti-H_CD40 hIgG1 Antibody(APX005M)
Anti-H_CD40 hIgG1 Antibody(ravagalimab)	Anti-H_CD40L hIgG1 Antibody(dapirolizumab)
Anti-H_CD40L hIgG1 Antibody(frexalimab)	
Biotinylated Human CD40 Protein; His-Avi Tag	Cynomolgus CD40 Protein; His Tag
Human CD40 Protein; His Tag	Human CD40L Protein; His Tag
IFN- α	
IFNα Reporter HEK-293 Cell Line	IFNα Reporter MDCK Cell Line
IFNα Reporter THP1 Cell Line	
BCMA:BAFFR:TACI	
H_BAFFR Jurkat Blockade Reporter Cell Line	H_BAFFR Reporter Cell Line
H_BCMA Reporter Cell Line	H_TACI Reporter Cell Line
Cynomolgus_BCMA CHO-K1 Cell Line	Cynomolgus_BCMA HEK-293 Cell Line
H_BAFFR CHO-K1 Cell Line	H_BCMA CHO-K1 Cell Line

H_BCMA HEK-293 Cell Line	Membrane Bound H_APRIL(Trimer) HEK-293 Cell Line
Anti-BAFF hIgG1 Antibody(belumumab)	Anti-BAFFR hIgG1 Antibody(ianalumab)
Anti-BCMA hIgG1 Antibody(Belantamab)	Anti-BCMA hIgG1 Antibody(SEA-BCMA)
Anti-BCMA hIgG4 Antibody(BCMB69)	Anti-CD3E×BCMA hIgG4 Reference Antibody (Tecbio)
Anti-TNFSF13B(BAFF) hIgG1 Reference Antibody (Belibio)	
Biotinylated Human BAFF Protein; His-Avi Tag	Biotinylated Human BCMA Protein; His-Avi Tag
Cynomolgus BAFF Protein; His Tag	Cynomolgus BCMA Protein; hFc Tag
Cynomolgus BCMA Protein; His Tag	Human APRIL Protein; hFc Tag
Human BAFF Protein; His Tag	Human BCMA Protein; hFc Tag
Human BCMA Protein; His Tag	Mouse BAFF Protein; His Tag
BDCA2(CLEC4C)	
H_BDCA2 Reporter DDX35TM Jurkat Cell Line	H_BDCA2 Reporter Jurkat Cell Line
Cynomolgus_BDCA2 CHO-K1 Cell Line	Cynomolgus_BDCA2 Jurkat Cell Line
H_BDCA2 CHO-K1 Cell Line	H_BDCA2 HEK-293 Cell Line
H_BDCA2 Jurkat Cell Line	
Anti-H_BDCA2 hIgG1 Antibody(Litifilimab)	
Cynomolgus BDCA2 Protein; His Tag	Human BDCA2 Protein; His Tag

License Agreement:

By purchasing and using this cell line product, the user voluntarily agrees to accept and abide by the following policies:

- This cell line product is restricted to research use only and shall not be used for any commercial purposes.
- This product is strictly prohibited from being used in the diagnosis or treatment of human or animal diseases, and shall not be directly used in experiments involving humans.
- Users and their contractors engaged for their benefit may use this material and its derivatives only within the agreed research scope; modification of the material is not permitted, nor may it be distributed, sold, transferred, or otherwise provided to any other entity (including affiliates).
- If use beyond the above scope is required, prior written permission from Genomeditech (Shanghai) Co.,Ltd. must be obtained. For details, please contact Genomeditech (Shanghai) Co.,Ltd.