Genomeditech (Shanghai) Co., Ltd.
Order: +86 021-50432825/50432826
Email: service@genomeditech.com

Product Sheet

H_CLDN18.2 HT29 Cell Line

Catalog number: GM-C48102

Version 4.0.26.1029

Specifications

Description	H_CLDN18.2 HT29 Cell Line is a clonal stable HT29 cell line that constitutively expresses the human CLDN18.2 gene, constructed using lentiviral technology.
Quantity	5E6 Cells per vial,1 mL
Product Format	1 vial of frozen cells
Shipping	Shipped on dry ice
Storage Conditions	Liquid nitrogen immediately upon receipt
Target	H_CLDN18.2
Gene ID/Uniprot ID	NP_001002026.1
Host Cell	HT29
Recovery Medium	McCoy's 5A+10% FBS+1% P.S
Growth medium	McCoy's 5A+10% FBS+1% P.S+0.5 μg/mL Puromycin
Note	None
Freezing Medium	90% FBS+10% DMSO
Growth properties	Adherent
Growth Conditions	37°C, 5% CO_2
Mycoplasma Testing	The cell line has been screened to confirm the absence of Mycoplasma species.
Safety considerations	Biosafety Level 2
Note	It is recommended to expand the cell culture and store a minimum of 10 vials at an early passage for potential future use.

Genomeditech

Materials

Reagent	Manufacturer/Catalogue No.
McCoy's 5A	VivaCell/C3020-0500
Fetal Bovine Serum	ExCell/FSP500
Pen/Strep	Thermo/15140-122
Puromycin	Genomeditech/GM-040401
Countess 3 Automated Cell Counter	ThermoFisher/A49891
NovoCyte Flow Cytometer	Agilent/NovoCyte 2060R
Anti-CLDN18.2 hIgG1 Antibody(LM-102)	Genomeditech/GM-81801AB

Data

Expression



	SampleID	Geometric Mean : R675-H
	HT29 Anti-CLDN18.2+APC 2nd Ab	1302
	HT29 H_CLDN18.2 H_IgG+APC 2nd Ab	1489
	HT29 H_CLDN18.2 Anti-CLDN18.2+APC 2nd Ab	53755

H_CLDN18.2 HT29 Cell Line (Cat. GM-C48102) was determined by flow cytometry using Anti-CLDN18.2 hIgG1 Antibody(LM-102)(Cat. GM-81801AB).

Cell Recovery

Recovery Medium: McCoy's 5A+10% FBS+1% P.S

To insure the highest level of viability, thaw the vial and initiate the culture as soon as possible upon receipt. If upon arrival, continued storage of the frozen culture is necessary, it should be stored in liquid nitrogen vapor phase and not at -70°C. Storage at -70°C will result in loss of viability.

a) Thaw the vial by gentle agitation in a 37°C water bath. To reduce the possibility of contamination, keep the O-ring and cap out of the water. Thawing should be rapid (approximately 2 - 3 minutes).
b) Remove the vial from the water bath as soon as the contents are thawed, and decontaminate by dipping in or spraying with 70% ethanol. All of the operations from this point on should be carried out under strict aseptic conditions.
c) Transfer the vial contents to a centrifuge tube containing 5.0 mL complete culture medium. And spin at approximately 176 x g for 5 minutes. Discard supernatant.
d) Adjust the viable cell density to 4–5E5 cells/mL by supplementing with recovery medium. Based on the total volume of the cell suspension, seed the cells into appropriate culture vessels.
e) Incubate the culture at 37°C in a suitable incubator. A 5% CO_2 in air atmosphere is recommended if using the medium described on this product sheet.

Cell Freezing

Freezing Medium: 90% FBS+10% DMSO

a) Centrifuge at 176 x g for 3 minutes to collect cells.
b) Resuspend the cells in pre-cooled freezing medium and adjust the cell density to 5E6 cells/mL.
c) Aliquot 1 mL into each vial.
d) Place the vial in a controlled-rate freezing container and store at -80°C for at least 1 day, then transfer to liquid nitrogen as soon as possible.

Cell passage

Growth medium: McCoy's 5A+10% FBS+1% P.S+0.5 μg/mL Puromycin

For the first 1 to 2 passages post-resuscitation, use the recovery medium. Once the cells have stabilized, switch to a growth medium.

a) It is recommended to incubate the cells in a humidified incubator for 36-48 hours before microscopic examination. It is important to maintain confluency below 80%, exceeding this density may result in contact inhibition-induced reductions in cell viability due to mechanical compression.
b) Remove and discard culture medium.
c) Briefly rinse the cell layer with PBS to remove all traces of serum that contains trypsin inhibitor.
d) Add 1.0 mL of 0.25% (w/v) Trypsin-EDTA solution to dish and observe cells under an inverted microscope until cell layer is dispersed (usually within 3 to 5 minutes at 37°C).
e) Note: To avoid clumping do not agitate the cells by hitting or shaking the flask while waiting for the cells to detach. Cells that are difficult to detach may be placed at 37°C to facilitate dispersal.
f) Add 2.0 mL of growth medium to mix well and aspirate cells by gently pipetting.
g) After centrifugation, resuspend the pellet and add appropriate aliquots of the cell suspension to new culture vessels.Incubate cultures at 37°C.(calculate seeding density according to the culture surface area and cell density to achieve a post-passage confluency of 25–35%).

h) Incubate cultures at 37°C.

Subcultivation Ratio: A subcultivation ratio of 1:2 - 1:3 is recommended

Medium Renewal: Every 3 to 4 days

Notes

a) FBS requires heat inactivation at 56°C for 30 minutes, which can inactivate complement and some viruses, but does not significantly affect the activity of most growth factors and cytokines.

b) HT29 cells grow in patch-like aggregates, and the presence of numerous dark granules in the cell body is a normal phenomenon.

c) During the first week after recovery, HT29 cells may show a large number of floating cells. Fresh culture medium can be replaced, and the floating cells should be collected by centrifugation and returned to the culture flask to be co-cultured with the adherent cells. Newly recovered cells grow in a patch-like pattern with three-dimensional contact growth (i.e., not as a monolayer). After cell division and proliferation, the cells will revert to an epithelial morphology and grow as a flat monolayer.

Related Products

CLDN18	
Cynomolgus_CLDN18.2-eGFP CHO-K1 Cell Line	H_CLDN18(isoform2)-eGFP 293 Cell Line
H_CLDN18.1-eGFP HEK-293 Cell Line	H_CLDN18.2 MC38 Cell Line
H_CLDN18.2 MKN45 Cell Line(High Expression)	H_CLDN18.2 MKN45 Cell Line(Low Expression)
H_CLDN18.2 MKN45 Cell Line(Medium Expression)	H_CLDN18.2 PANC-1 Cell Line
H_CLDN18.2(isoform2) CHO-K1 Cell Line	H_CLDN18.2-eGFP CT-26 Cell Line
Mouse_CLDN18.2-eGFP CHO-K1 Cell Line	Rat_CLDN18.2-eGFP CHO-K1 Cell Line
Rhesus_CLDN18.2-eGFP CHO-K1 Cell Line	
Anti-CLDN18.2 hIgG1 Reference Antibody (IMAB362)	Anti-CLDN18.2 hIgG1 Antibody(LM-102)
Anti-CLDN18.2 hIgG1 Antibody(Zolbetuximab)	
TROP2(TACSTD2)	
Cynomolgus_Trop2 CHO-K1 Cell Line	Cynomolgus_TROP2 HEK-293 Cell Line
H_TROP2 CHO-K1 Cell Line	H_TROP2 CT26 Cell Line
H_TROP2 HEK-293 Cell Line	H_TROP2 LLC1 Cell Line
Anti-H_TROP2 hIgG1 Antibody(Datopotamab)	Anti-TROP2 hIgG1 Antibody(Hu2G10-5)
Anti-Trop2 hIgG1 Reference Antibody (Sacbio)	Anti-Trop2 hIgG1 Reference Antibody(Datbio)
Anti-Trop2-DXD ADC(Dar4)[Datopotamab deruxtecan,Dato-DXD]	Anti-Trop2-SN38 ADC(Dar8)[Sacituzumab govitecan]
Biotinylated Cynomolgus TROP2 Protein; His-Avi Tag	Biotinylated Human TROP2 Protein; His-Avi Tag
Human TROP2 Protein; hFc Tag	Human TROP2 Protein; His Tag
GUCY2C(GC-C)	
Cynomolgus_GUCY2C HEK-293 Cell Line	H_GUCY2C CHO-K1 Cell Line
H_GUCY2C HCT116 Cell Line	H_GUCY2C HEK-293 Cell Line
Anti-H_GUCY2C hIgG1 Antibody(Indusatumab)	
Cynomolgus GUCY2C Protein; His Tag	Human GUCY2C Protein; His Tag

CD3	
Cynomolgus_CD3D CD3E Reporter Jurkat(CD3DE KO) Cell Line	Jurkat CD3-BsAb Reporter Cell Line
Cynomolgus_CD3 HEK-293 Cell Line	Cynomolgus_CD3E(Membrane Bound ECD) CHO-K1 Cell Line
H_CD3 CHO-K1 Cell Line	H_CD3 HEK-293 Cell Line
H_CD3(TCR V2) CHO-K1 Cell Line	H_CD3(TCR V2) HEK-293 Cell Line
H_CD3(TCRαβ delta V) HEK-293 Cell Line	H_CD3D CD3E KO Jurkat Cell Line
H_CD3E KO Jurkat Cell Line	H_CD3E(Membrane Bound ECD) CHO-K1 Cell Line
Mouse_CD3 HEK-293 Cell Line	
Anti-CD19×CD3 hIgG1 Antibody[PIT-565(CD58 K34A)]	Anti-CD3 epsilon hIgG1 Antibody [OKT-3 (muromonab)]
Anti-CD3 hIgG1 Antibody(CH2527)	Anti-CD3×CD20 hIgG1 Bispecific Antibody (Epcobio)
Anti-CD3×FCRL5 hIgG1 Bispecific Antibody(cevostamab)	Anti-CD3E×BCMA hIgG4 Reference Antibody (Tecbio)
Anti-CD3E×DLL3 hIgG1 Bispecific Antibody(Tarlabio)	Anti-CD3E×MUC17 hIgG1 Bispecific Antibody(Vepsitbio)
Anti-mouse CD3ε mIgG2a Antibody(145-2C11)	
Cynomolgus CD3E Protein; hFc Tag	Human CD3E Protein; hFc Tag
FGFR2	
H_FGFR2b Reporter 293 Cell Line	Cynomolgus_FGFR2b HEK-293 Cell Line
H_FGFR2b CHO-K1 Cell Line	H_FGFR2b HEK-293 Cell Line
Mouse_FGFR2b HEK-293 Cell Line	
Anti-FGFR2 hIgG1 Antibody(hFR2-14_H12-L1)	Anti-FGFR2 hIgG1 Referece Antibody (Bemabio)
Cynomolgus FGFR2(IIIb) D1-D3 Protein; His Tag	Human FGFR2(IIIb) D1-D3 Protein; hFc Tag
Human FGFR2(IIIb) D2-D3 Protein; hFc Tag	Human FGFR2(IIIb) D2-D3 Protein; His Tag
Human FGFR2(IIIc) D1-D3 Protein; His Tag	Mouse FGFR2(IIIb) D2-D3 Protein; His Tag
CLDN3	
H_CLDN3 HEK-293 Cell Line	
Anti-CLDN3 hIgG1 Antibody(H4G3)	
CLDN4	
H_CLDN4 HEK-293 Cell Line	
Anti-CLDN4 hIgG1 Antibody(4B8)	
CLDN6	
Cynomolgus_CLDN6 CHO-K1 Cell Line	H_CLDN6 CHO-K1 Cell Line
H_CLDN6 HEK-293 Cell Line	H_CLDN6 LLC1 Cell Line
H_CLDN6 MC38 Cell Line	Mouse_CLDN6 CHO-K1 Cell Line
Rat_CLDN6 CHO-K1 Cell Line	
Anti-Claudin6 hIgG1 Reference Antibody	Anti-CLDN6/9 hIgG1 Antibody
CLDN9	
H_CLDN9 CHO-K1 Cell Line	H_CLDN9-eGFP HEK-293 Cell Line
CLDN1	
Cynomolgus_CLDN1 CHO-K1 Cell Line	H_CLDN1 CHO-K1 Cell Line
H_CLDN1 HCT116 Cell Line	Mouse_CLDN1-GFP CHO-K1 Cell Line
Anti-CLDN1 hIgG1 Reference Antibody (Lixubio)	

CD44	
H_CD44s CHO-K1 Cell Line	H_CD44v6 CHO-K1 Cell Line
H_CD44v6 CHO-K1 Cell Line (Medium Expression)	H_CD44v6 HEK-293 Cell Line
Anti-CD44v6 hIgG1 Antibody(bivatuzumab)	
ADC Related Product	
Anti-DXD Mouse IgG1 Antibody (23E21C5)	Anti-DXD Mouse IgG1 Antibody (4A5A12)
Anti-Dxd Mouse IgG2a Antibody (17D6A4)	Anti-Eribulin Mouse IgG2a Antibody (10F8G4)
Anti-MMAE Mouse IgG1 Antibody (11C10E3)	Anti-MMAE Mouse IgG2a Antibody (17A1K11)
Anti-MMAE Mouse IgG2a Antibody (8F6A3)	Anti-SN38 Mouse IgG1 Antibody(59H11C7)
Mouse anti Human IgG1-DXD(Dar8)	Mouse anti Human IgG1-MMAE(Dar4)
Human IgG1 Isotype-DXD (Dar8)	Human IgG1 Isotype-Eribulin (Dar4)
Human IgG1 Isotype-MMAE (Dar4)	
Recombinant DT3C Protein	

License Agreement

By purchasing and using this cell line product, the user voluntarily agrees to accept and abide by the following policies:

- This cell line product is restricted to research use only and shall not be used for any commercial purposes.
- This product is strictly prohibited from being used in the diagnosis or treatment of human or animal diseases, and shall not be directly used in experiments involving humans.
- Users and their contractors engaged for their benefit may use this material and its derivatives only within the agreed research scope; modification of the material is not permitted, nor may it be distributed, sold, transferred, or otherwise provided to any other entity (including affiliates).
- If use beyond the above scope is required, prior written permission from Genomeditech (Shanghai) Co.,Ltd. must be obtained. For details, please contact Genomeditech (Shanghai) Co.,Ltd.

Genomeditech