

Product Sheet

IL-3 Reporter 293 Cell Line


Catalog number: GM-C26142

Version 3.3.1.260603

IL-3 (Interleukin-3) is a cytokine produced by T cells, mast cells, and monocytes, playing a crucial role in hematopoiesis and immune responses. It stimulates the proliferation and differentiation of hematopoietic stem cells, promoting the generation of red blood cells, white blood cells, and platelets, while also enhancing immune cell functions against infections and tumors.

The IL-3 signaling pathway is mediated by the IL-3R(IL-3 α and β chain). When IL-3 binds to IL-3R, it activates JAK2 and other tyrosine kinases, which then activate transcription factors like STAT5. These factors enter the nucleus to regulate target gene expression, promoting hematopoietic cell proliferation and differentiation. Abnormalities in this pathway may be linked to hematological diseases such as leukemia and lymphoma.

IL-3 Reporter 293 Cell Line is a clonal stable HEK-293 cell line constructed using lentiviral technology, constitutive expression of the IL-3 α and β chain, along with signal-dependent expression of a luciferase reporter gene. When IL-3 binds to IL-3R, it activates downstream signaling pathways, leading to the expression of luciferase. Blockade antibodies can inhibit this signal transmission. The luciferase activity measurement indicates the activation level of the signaling pathway and can thus be used to evaluate the in vitro effects of drugs related to IL-3.



Specifications

Quantity	5E6 Cells per vial, 1 mL
Product Format	1 vial of frozen cells
Shipping	Shipped on dry ice
Storage Conditions	Liquid nitrogen immediately upon receipt
Recovery Medium	DMEM+10% FBS+1% P.S
Growth medium	DMEM+10% FBS+1% P.S+4 µg/mL Blasticidin+125 µg/mL Hygromycin+0.75 µg/mL Puromycin
Note	None
Freezing Medium	90% FBS+10% DMSO
Growth properties	Adherent
Growth Conditions	37°C, 5% CO ₂
Mycoplasma Testing	The cell line has been screened to confirm the absence of Mycoplasma species.
Safety considerations	Biosafety Level 2
Note	It is recommended to expand the cell culture and store a minimum of 10 vials at an early passage for potential future use.

Materials

Reagent	Manufacturer/Catalogue No.
DMEM	Gibco/C11995500BT
Fetal Bovine Serum	ExCell/FSP500
Pen/Strep	Thermo/15140-122
Blasticidin	Genomeditech/ GM-040404
Hygromycin	Genomeditech/ GM-040403
Puromycin	Genomeditech/ GM-040401
Recombinant Human IL-3 (C-6His)	Novoprotein/CX90
Anti-H ₂ CD123 hIgG2 Antibody(Talacotuzumab)	Genomeditech/ GM-27123AB
GMOne-Step 2.0 Luciferase Reporter Gene Assay Kit	Genomeditech/ GM-040513

Figures




Figure 1 | Response to Recombinant Human IL-3 (C-6His). The IL-3 Reporter 293 Cell Line (Cat. GM-C26142) at a concentration of 1.5E4 cells/well (96-well format) was stimulated with serial dilutions of Recombinant Human IL-3 (C-6His) (Novoprotein/CX90) in assay buffer (DMEM + 1% FBS + 1% P.S) for 7 hours. The firefly luciferase activity was measured using the Luciferase Reporter Assay Kit (Genomeditech). The maximum induction fold was approximately [20.1]. Data are shown by drug mass concentration.




Figure 2 | Response to Anti-H_CD123 hIgG2 Antibody(Talacotuzumab). Serial dilutions of the Anti-H_CD123 hIgG2 Antibody(Talacotuzumab) (Cat. [GM-27123AB](#)) was incubated with 1.5E4 cells/well of the IL-3 Reporter 293 Cell Line (Cat. GM-C26142) in a 96-well plate for 1 hour in assay buffer (DMEM + 1% FBS + 1% P.S). Subsequently, the Human Recombinant Human IL-3 (C-6His) (Novoprotein/CX90) at a concentration of 0.4 ng/well was added, and the coculture proceeded for an additional 6 hours. Firefly luciferase activity is then measured using the Luciferase Reporter Assay Kit (Genomeditech). The results indicated maximum blocking folds of approximately [17.7]. Data are shown by drug mass concentration.




Figure 3 | Response to Anti-H_CD123 hIgG2 Antibody(Talacotuzumab). Serial dilutions of the Anti-H_CD123 hIgG2 Antibody(Talacotuzumab) (Cat. [GM-27123AB](#)) was incubated with 1.5E4 cells/well of the IL-3 Reporter 293 Cell Line (Cat. GM-C26142) in a 96-well plate for 1 hour in assay buffer (DMEM + 1% FBS + 1% P.S). Subsequently, the Human Recombinant Human IL-3 (C-6His) (Novoprotein/CX90) at a concentration of 0.4 ng/well was added, and the coculture proceeded for an additional 6 hours. Firefly luciferase activity is then measured using the Luciferase Reporter Assay Kit (Genomeditech). The results indicated maximum blocking folds of approximately [17.7]. Data are shown by drug molar concentration.




Figure 4 | IL-3 Reporter 293 Cell Line (Cat. GM-C26142) was determined by flow cytometry using Anti-H_CD123 hIgG2 Antibody(Talacotuzumab) (Cat. [GM-27123AB](#)).

Cell Recovery

Recovery Medium: DMEM+10% FBS+1% P.S

To insure the highest level of viability, thaw the vial and initiate the culture as soon as possible upon receipt. If upon arrival, continued storage of the frozen culture is necessary, it should be stored in liquid nitrogen vapor phase and not at -70°C . Storage at -70°C will result in loss of viability.

- Thaw the vial by gentle agitation in a 37°C water bath. To reduce the possibility of contamination, keep the O-ring and cap out of the water. Thawing should be rapid (approximately 2 - 3 minutes).
- Remove the vial from the water bath as soon as the contents are thawed, and decontaminate by dipping in or spraying with 70% ethanol. All of the operations from this point on should be carried out under strict aseptic conditions.
- Transfer the vial contents to a centrifuge tube containing 5.0 mL complete culture medium and spin at approximately $176 \times g$ for 5 minutes. Discard supernatant.
- Resuspend cell pellet with the recommended recovery medium. And dispense into appropriate culture dishes.
- Incubate the culture at 37°C in a suitable incubator. A 5% CO_2 in air atmosphere is recommended if using the medium described on this product sheet.

Cell Freezing

Freezing Medium: 90% FBS+10% DMSO

- Centrifuge at $176 \times g$ for 3 minutes to collect cells.
- Resuspend the cells in pre-cooled freezing medium and adjust the cell density to 5×10^6 cells/mL.
- Aliquot 1 mL into each vial.
- Place the vial in a controlled-rate freezing container and store at -80°C for at least 1 day, then transfer to liquid nitrogen as soon as possible.

Cell passage

Growth medium: DMEM+10% FBS+1% P.S+4 $\mu\text{g/mL}$ Blasticidin+125 $\mu\text{g/mL}$ Hygromycin+0.75 $\mu\text{g/mL}$ Puromycin

For the first 1 to 2 passages post-resuscitation, use the recovery medium. Once the cells have stabilized, switch to a growth medium.

- Subculturing is necessary when the cell density reaches 80%. It is recommended to perform subculturing at a ratio of 1:3 to 1:4 every 2-3 days. Ensure that the density does not exceed 80%, as overcrowding can lead to reduced viability due to compression.
- Remove and discard culture medium.
- Briefly rinse the cell layer with PBS to remove all traces of serum that contains trypsin inhibitor.
- Add 1.0 mL of 0.25% (w/v) Trypsin-EDTA solution to dish and observe cells under an inverted microscope until cell layer is dispersed (usually within 30 to 60 seconds at 37°C).
- Note: To avoid clumping do not agitate the cells by hitting or shaking the flask while waiting for the cells to detach. Cells that are difficult to detach may be placed at 37°C to facilitate dispersal.
- Add 2.0 mL of growth medium to mix well and aspirate cells by gently pipetting.
- After centrifugation, resuspend the pellet and add appropriate aliquots of the cell suspension to new culture vessels.
- Incubate cultures at 37°C .

Subcultivation Ratio: A subcultivation ratio of 1:3 - 1:4 is recommended

Medium Renewal: Every 2 to 3 days

Notes

- Upon initial thawing, a higher number of dead cells is observed, which is a normal phenomenon. Significant improvement is seen after adaptation. Once the cells reach a stable state, the number of dead cells decreases after subculturing and the cell growth rate becomes stable.
- Ensure that the cell density does not exceed 80%, as overcrowding may lead to reduced viability due to compression.

Related Products

IL-3	
Cynomolgus_CD123(IL3RA) CHO-K1 Cell Line	H_CD123(IL3RA) CHO-K1 Cell Line
H_CD123(IL3RA) HEK-293 Cell Line	
Anti-H_CD123 hIgG2 Antibody(Talacotuzumab)	

License Agreement:

By purchasing and using this cell line product, the user voluntarily agrees to accept and abide by the following policies:

- This cell line product is restricted to research use only and shall not be used for any commercial purposes.
- This product is strictly prohibited from being used in the diagnosis or treatment of human or animal diseases, and shall not be directly used in experiments involving humans.
- Users and their contractors engaged for their benefit may use this material and its derivatives only within the agreed research scope; modification of the material is not permitted, nor may it be distributed, sold, transferred, or otherwise provided to any other entity (including affiliates).
- If use beyond the above scope is required, prior written permission from Genomeditech (Shanghai) Co.,Ltd. must be obtained. For details, please contact Genomeditech (Shanghai) Co.,Ltd.