

# Product Sheet

## Mouse\_FcgRIV FcgRIIb CHO-K1 Cell Line

Catalog number: GM-C40208

Version 3.3.1.260624

<b>Description</b>	Mouse_FcgRIV FcgRIIb CHO-K1 Cell Line is a clonal stable CHO-K1 cell line that constitutively expresses the Mouse FcgRIV and FcgRIIb genes, constructed using lentiviral technology.
<b>Quantity</b>	5E6 Cells per vial, 1 mL
<b>Product Format</b>	1 vial of frozen cells
<b>Shipping</b>	Shipped on dry ice
<b>Storage Conditions</b>	Liquid nitrogen immediately upon receipt
<b>Target</b>	Mouse_FcgRIV FcgRIIb
<b>Gene ID/Uniprot ID</b>	P08101-1 1997-11-01 v2、A0A0B4J1G0 2015-03-04 v1
<b>Host Cell</b>	CHO-K1
<b>Recovery Medium</b>	F12K+10% FBS+1% P.S
<b>Growth medium</b>	F12K+10% FBS+1% P.S+100 µg/mL Hygromycin+4 µg/mL Puromycin
<b>Note</b>	None
<b>Freezing Medium</b>	90% FBS+10% DMSO
<b>Growth properties</b>	Adherent
<b>Growth Conditions</b>	37°C, 5% CO <sub>2</sub>
<b>Mycoplasma Testing</b>	The cell line has been screened to confirm the absence of Mycoplasma species.
<b>Safety considerations</b>	Biosafety Level 2
<b>Note</b>	It is recommended to expand the cell culture and store a minimum of 10 vials at an early passage for potential future use.

## Materials

Reagent	Manufacturer/Catalogue No.
Puromycin	Genomeditech/GM-040401
Blasticidin	Genomeditech/GM-040404
Hygromycin	Genomeditech/GM-040403
Pen/Strep	Thermo/15140-122
Fetal Bovine Serum	ExCell/FSP500
F12K	BOSTER/PYG0036
APC anti-mouse CD16.2 (FcγRIV) Antibody	Biolegend/149505
FITC anti-mouse CD32 (FcγR2)	Biolegend/156407
Anti-TREM1 mIgG1 Antibody(PY159)	Genomeditech/GM-88439AB
Anti-TREM1 mIgG2a Antibody(PY159)	Genomeditech/GM-88440AB
Anti-TREM1 mIgG2b Antibody(PY159)	Genomeditech/GM-88442AB
H_TREM1 Reporter Jurkat Cell Line	Genomeditech/GM-C44546

## Figures




Figure 1 | Response to Anti-TREM1 mIgG2a Antibody(PY159). The Mouse\_FcgRIV FcgRIIb CHO-K1 Cell Line (Cat. GM-C40208) was seeded at a density of 1E4 cells per well in a 96-well plate and incubated overnight. The next day, serial dilutions of Anti-TREM1 mIgG2a Antibody (PY159, Cat. GM-88440AB) and the H\_TREM1 Reporter Jurkat Cell Line (Cat. GM-C44546) at a density of 1E5 cells per well were added to the pre-seeded cells. The mixture was incubated for an additional 6 hours. Firefly luciferase activity was measured using the Luciferase Reporter Assay Kit (Genomeditech). The maximum induction fold was approximately [179.9]. Data are presented based on drug mass concentration.




Figure 2 | Mouse\_FcgRIV FcgRIIb CHO-K1 Cell Line (Cat. GM-C40208) was determined by flow cytometry using APC anti-mouse CD16.2 (FcyRIV) Antibody (Biolegend/149505).




Figure 3 | Mouse\_FcgRIV FcgRIIb CHO-K1 Cell Line (Cat. GM-C40208) was determined by flow cytometry using FITC anti-mouse CD32 (Fcgr2) (Biolegend/156407).

## Cell Recovery

Recovery Medium: F12K+10% FBS+1% P.S

To insure the highest level of viability, thaw the vial and initiate the culture as soon as possible upon receipt. If upon arrival, continued storage of the frozen culture is necessary, it should be stored in liquid nitrogen vapor phase and not at -70°C. Storage at -70°C will result in loss of viability.

- Thaw the vial by gentle agitation in a 37°C water bath. To reduce the possibility of contamination, keep the O-ring and cap out of the water. Thawing should be rapid (approximately 2 - 3 minutes).
- Remove the vial from the water bath as soon as the contents are thawed, and decontaminate by dipping in or spraying with 70% ethanol. All of the operations from this point on should be carried out under strict aseptic conditions.
- Transfer the vial contents to a centrifuge tube containing 5.0 mL complete culture medium and spin at approximately 176 x g for 5 minutes. Discard supernatant.
- Resuspend cell pellet with the recommended recovery medium. And dispense into appropriate culture dishes.

- e) Incubate the culture at 37°C in a suitable incubator. A 5% CO<sub>2</sub> in air atmosphere is recommended if using the medium described on this product sheet.

## Cell Freezing

Freezing Medium: 90% FBS+10% DMSO

- Centrifuge at 176 x g for 3 minutes to collect cells.
- Resuspend the cells in pre-cooled freezing medium and adjust the cell density to 5E6 cells/mL.
- Aliquot 1 mL into each vial.
- Place the vial in a controlled-rate freezing container and store at -80°C for at least 1 day, then transfer to liquid nitrogen as soon as possible.

## Cell passage

Growth medium: F12K+10% FBS+1% P.S+100 µg/mL Hygromycin+4 µg/mL Puromycin

For the first 1 to 2 passages post-resuscitation, use the recovery medium. Once the cells have stabilized, switch to a growth medium.

- Remove and discard culture medium.
- Briefly rinse the cell layer with PBS to remove all traces of serum that contains trypsin inhibitor.
- Add 1.0 mL of 0.25% (w/v) Trypsin-EDTA solution to dish and observe cells under an inverted microscope until cell layer is dispersed (usually within 2 to 3 minutes at 37°C).
- Note: To avoid clumping do not agitate the cells by hitting or shaking the flask while waiting for the cells to detach. Cells that are difficult to detach may be placed at 37°C to facilitate dispersal.
- Add 2.0 mL of growth medium to mix well and aspirate cells by gently pipetting.
- After centrifugation, resuspend the pellet and add appropriate aliquots of the cell suspension to new culture vessels.
- Incubate cultures at 37°C.

**Subcultivation Ratio: A subcultivation ratio of 1:4 - 1:5 is recommended**

**Medium Renewal: Every 2 to 3 days**

## Notes

- After the stabilization of the cell condition, there will be fewer dead cells post-passage, the cell growth rate will tend to stabilize, cell morphology will become uniform, and the cells will appear robust.

## Sequence

Fcgr2b [P08101-1](#)

MESNWTVHVFSRTLCHMLLWTAVLNLAAGTHDLPKAVVKLEPPWIQVLKEDTVTLTCEGTHNPGNSSTQW  
FHNGRSIRSQVQASYTFKATVNDSGEYRCQMEQTRLSDPVDLGVISDWLLLQTPQLVFLEGETITLRCHSWR  
NKLLNRISFFHNEKSVRYHHYSSNFSIPKANHSHSGDYCYCKGSLGRTLHQSKPVTITVQGPKSSRSLPVLTVIA

AVTGIAVA AIVILVSLVYLKKKQVPALPGNPDHREMGETLPEEVGEYRQPSGGSVPVSPGPPSGLEPTSSSPY  
 NPPDLEEA AKTEA ENTITYSLLKHPEALDEETEHDYQNH I

FcgRIV [A0A0B4J1G0](#)

MWQLLLPTALVLTAFSGIQAGLQKAVVNLDPKWVRVLEEDSVTLRCQGTFSPEDNSIKWFHNESLIPHQDAN  
 YVIQSARVKDSGMYRCQTALSTISDPVQLEVHMGWLLLQTTKWLFQEGDPIHLRCHSWQNRVPRKVTYLQN  
 GKGKKYFHENSELLIPKATHNDSGSYFCRGLIGHNNKSSASFRISLGDGPGSPSMFPPWHQITFCLLIGLLFAIDT  
 VLYFSVRRGLQSPVADYEEP KIQWSKEPQDK

## Related Products

FcγR	
<a href="#">H_CD32B aAPC CHO-K1 Cell Line</a>	<a href="#">Cynomolgus_FcRn MDCK Cell Line</a>
<a href="#">H_FCGR1A(CD64) CHO-K1 Cell Line</a>	<a href="#">H_FCGR1A(CD64) HEK-293 Cell Line</a>
<a href="#">H_FCGR2A(CD32A) CHO-K1 Cell Line</a>	<a href="#">H_FCGR2B(CD32B) CHO-K1 Cell Line</a>
<a href="#">H_FCGR3A(CD16a) 158F CHO-K1 Cell Line</a>	<a href="#">H_FCGR3A(CD16a) 158V CHO-K1 Cell Line</a>
<a href="#">H_FCGR3B(CD16b) CHO-K1 Cell Line</a>	<a href="#">H_FcRn CHO-K1 Cell Line</a>
<a href="#">H_FcRn MDCK Cell Line</a>	<a href="#">H_FcRn-GFP HEK-293 Cell Line</a>
<a href="#">Mouse_FcgRIV FcgRIIb aAPC CHO-K1 Cell Line</a>	<a href="#">Mouse_FcRn MDCK Cell Line</a>
<a href="#">Anti-FcRn hIgG4 Reference Antibody(Rozabio)</a>	<a href="#">Anti-H_FcRn IgG4 Antibody(Rozanolixizumab)</a>
<a href="#">Anti-Mouse CD16/32 mIgG2b Antibody(2.4G2)</a>	
<a href="#">Human IgG1 Fc Protein; His Tag</a>	
ADCCP	
<a href="#">ADCC FcγRIIIa(158F) Jurkat Effector Cell Line</a>	<a href="#">ADCC FcγRIIIa(158V) DDX35TM Jurkat Effector Cell Line</a>
<a href="#">ADCC FcγRIIIa(158V) Jurkat Effector Cell Line</a>	<a href="#">ADCC FcγRIIIa(158V) Reporter Jurkat(CD3 KO) Cell Line</a>
<a href="#">ADCC FcγRIIIa(158V) Reporter Jurkat(CD3 KO) Cell Line</a>	<a href="#">ADCC FcγRIIIa(158V) Reporter NK-92 Cell Line</a>
<a href="#">ADCC M_FcγRIV Jurkat Effector Cell Line</a>	<a href="#">ADCP FcγRI Jurkat Effector Cell Line</a>
<a href="#">ADCP FcγRIIa DDX35TM Jurkat Effector Cell Line</a>	<a href="#">ADCP FcγRIIa Jurkat Effector Cell Line</a>
<a href="#">ADCP FcγRIIa R131 Jurkat Effector Cell Line</a>	<a href="#">ADCP FcγRIIb Jurkat Effector Cell Line</a>

## License Agreement:

**By purchasing and using this cell line product, the user voluntarily agrees to accept and abide by the following policies:**

- This cell line product is restricted to research use only and shall not be used for any commercial purposes.
- This product is strictly prohibited from being used in the diagnosis or treatment of human or animal diseases, and shall not be directly used in experiments involving humans.
- Users and their contractors engaged for their benefit may use this material and its derivatives only within the agreed research scope; modification of the material is not permitted, nor may it be distributed, sold, transferred, or otherwise provided to any other entity (including affiliates).

- If use beyond the above scope is required, prior written permission from Genomeditech (Shanghai) Co.,Ltd. must be obtained. For details, please contact Genomeditech (Shanghai) Co.,Ltd.

Genomeditech