

Product Sheet

Raji Cell Line

Catalog number: GM-C19100

Version 3.3.1.260723

Description	Raji cells are used as the feeder cell line for H_CTLA4 Reporter Jurkat Cells (Cat. GM-C23902) and constitutively express CD80.
Quantity	5E6 Cells per vial, 1 mL
Product Format	1 vial of frozen cells
Shipping	Shipped on dry ice
Storage Conditions	Liquid nitrogen immediately upon receipt
Target	/
Gene ID/Uniprot ID	/
Host Cell	Raji
Recovery Medium	RPMI 1640+10% FBS+1% P.S
Growth medium	RPMI 1640+10% FBS+1% P.S
Note	None
Freezing Medium	90% FBS+10% DMSO
Growth properties	Suspension
Growth Conditions	37°C, 5% CO ₂
Mycoplasma Testing	The cell line has been screened to confirm the absence of Mycoplasma species.
Safety considerations	Biosafety Level 2
Note	It is recommended to expand the cell culture and store a minimum of 10 vials at an early passage for potential future use.

Materials

Reagent	Manufacturer/Catalogue No.
RPMI 1640	VivaCell/C3010-0500
Fetal Bovine Serum	ExCell/FSP500
Pen/Strep	Thermo/15140-122
Anti-H_CD80 hIgG1 Antibody(Galiximab)	Genomeditech/ GM-46075AB

Figures




Figure 1 | Raji Cell Line (Cat. GM-C19100) was determined by flow cytometry using Anti-H_CD80 hIgG1 Antibody(Galiximab) (Cat. [GM-46075AB](#)).

Cell Recovery

Recovery Medium: RPMI 1640+10% FBS+1% P.S

To insure the highest level of viability, thaw the vial and initiate the culture as soon as possible upon receipt. If upon arrival, continued storage of the frozen culture is necessary, it should be stored in liquid nitrogen vapor phase and not at -70°C. Storage at -70°C will result in loss of viability.

- Thaw the vial by gentle agitation in a 37°C water bath. To reduce the possibility of contamination, keep the O-ring and cap out of the water. Thawing should be rapid (approximately 2 - 3 minutes).
- Remove the vial from the water bath as soon as the contents are thawed, and decontaminate by dipping in or spraying with 70% ethanol. All of the operations from this point on should be carried out under strict aseptic conditions.
- Transfer the vial contents to a centrifuge tube containing 5.0 mL complete culture medium. And spin at approximately 176 x g for 5 minutes. Discard supernatant.

- d) Resuspend cell pellet with the recommended complete medium. And dispense the suspension into an appropriate culture flask and initially place the flask in an upright position after thawing.
- e) Incubate the culture at 37°C in a suitable incubator. A 5% CO₂ in air atmosphere is recommended if using the medium described on this product sheet.

Cell Freezing

Freezing Medium: 90% FBS+10% DMSO

- a) Centrifuge at 176 x g for 3 minutes to collect cells.
- b) Resuspend the cells in pre-cooled freezing medium and adjust the cell density to 5E6 cells/mL.
- c) Aliquot 1 mL into each vial.
- d) Place the vial in a controlled-rate freezing container and store at -80°C for at least 1 day, then transfer to liquid nitrogen as soon as possible.

Cell passage

Growth medium: RPMI 1640+10% FBS+1% P.S

Antibiotics are not required during cell culture. The composition of the growth medium is identical to that of the recovery medium.

- a) When the cell density reaches 1 - 1.2E6 cells/mL, subculture the cells. Do not allow the cell density to exceed 1.2E6 cells/mL.
- b) It is recommended to use T-25 flasks for subculturing.
- c) These cells are suspension cells, and it is recommended to use the "half-medium change" method to maintain optimal cell conditions during passaging.
- d) During passaging, you can directly add fresh growth medium to the culture flask, gently pipette to resuspend the cells, and then transfer the cell suspension to a new T-25 flask for continued culture.

Subcultivation Ratio: Maintain cultures at a cell concentraion between 2.5E5 and 8E5 viable cells/mL.

Medium Renewal: Every 2 to 3 days

Notes

- a) FBS should be heat-inactivated at 56°C for 30 minutes to inactivate complement and certain viruses, with minimal impact on most growth factor and cytokine activities.

Related Products

CTLA4:CD80:CD86	
H_CD80 aAPC CHO-K1 Cell Line	H_CD80 PDL1 aAPC CHO-K1 Cell Line
H_CTLA4 PD-1 Reporter Cell Line	H_CTLA4 Reporter Jurkat Cell Line
H_CTLA4 Reporter(IL-2 promoter) Jurkat Cell Line	Canine_CTLA4 CHO-K1 Cell Line
Cynomolgus_CTLA4 HEK-293 Cell Line	H_CTLA4 (Full-Length) CHO-K1 Cell Line

H_CTLA4 CHO-K1 Cell Line	H_CTLA4 HEK-293 Cell Line
H_CTLA4 Jurkat Cell Line	H_CTLA4 PD-1 CHO-K1 Cell Line (High Ratio)
H_CTLA4 PD-1 CHO-K1 Cell Line (Low Ratio)	
Anti-CTLA4 hIgG1 Reference Antibody (Ipibio)	Anti-CTLA-4/PD-1 hIgG1 Bispecific Antibody(Cadonilimab)
Anti-CTLA4×PD-1 hIgG1 Reference Antibody (Cadbio)	Anti-H_CD80 hIgG1 Antibody(Galiximab)
Anti-H_CTLA-4 hIgG1 Antibody(Ipilimumab)	Anti-mouse CTLA4 mIgG2b Antibody(9D9)
Anti-mouse CTLA4 Syrian Hamster IgG2 Antibody(9H10)	
Biotinylated Cynomolgus CTLA4 Protein; His-Avi Tag	Biotinylated Mouse CTLA4 Protein; His-Avi Tag
Cynomolgus CTLA4 Protein; His Tag	Human CD80 Protein; His Tag
Human CTLA4 Protein; His Tag	Mouse CTLA4 Protein; His Tag

License Agreement:

By purchasing and using this cell line product, the user voluntarily agrees to accept and abide by the following policies:

- This cell line product is restricted to research use only and shall not be used for any commercial purposes.
- This product is strictly prohibited from being used in the diagnosis or treatment of human or animal diseases, and shall not be directly used in experiments involving humans.
- Users and their contractors engaged for their benefit may use this material and its derivatives only within the agreed research scope; modification of the material is not permitted, nor may it be distributed, sold, transferred, or otherwise provided to any other entity (including affiliates).
- If use beyond the above scope is required, prior written permission from Genomeditech (Shanghai) Co.,Ltd. must be obtained. For details, please contact Genomeditech (Shanghai) Co.,Ltd.