

# Product Sheet


## H\_CD3(TCR V2) HEK-293 Cell Line

Catalog number: GM-C44753

Version 3.3.1.260323

The CD3 complex, also known as the T3 complex, is a multimeric protein complex composed of four different polypeptide chains: epsilon ( $\epsilon$ ), gamma ( $\gamma$ ), delta ( $\delta$ ), and zeta ( $\zeta$ ). These chains assemble and function as three pairs of dimers:  $\epsilon\gamma$ ,  $\epsilon\delta$ , and  $\zeta\zeta$ . The CD3 protein complex is a defining feature of the T-cell lineage, thus anti-CD3 antibodies can be effectively used as markers for T cells. In T cells, the CD3 complex non-covalently associates with the T-cell receptor (TCR)  $\alpha$  and  $\beta$  chains to form the TCR-CD3 signaling complex, which is essential for antigen recognition and T-cell activation.

H\_CD3(TCR V2) HEK-293 Cell Line is a clonal stable HEK-293 cell line constitutively expressing human CD3 complex along with the T-cell receptor alpha and beta (TCR $\alpha$   $\beta$ ) chains. The key distinction between this cell line and the GM-C30877 H\_CD3 HEK-293 Cell Line lies in the sequence of the TCR components.



---

## Specifications

---

<b>Quantity</b>	5E6 Cells per vial, 1 mL
<b>Product Format</b>	1 vial of frozen cells
<b>Shipping</b>	Shipped on dry ice
<b>Storage Conditions</b>	Liquid nitrogen immediately upon receipt

---

<b>Recovery Medium</b>	DMEM+10% FBS+1% P.S
<b>Growth medium</b>	DMEM+10% FBS+1% P.S+400 µg/mL G418+125 µg/mL Hygromycin
<b>Note</b>	None
<b>Freezing Medium</b>	90% FBS+10% DMSO
<b>Growth properties</b>	Adherent
<b>Growth Conditions</b>	37°C, 5% CO <sub>2</sub>

---

<b>Mycoplasma Testing</b>	The cell line has been screened to confirm the absence of Mycoplasma species.
<b>Safety considerations</b>	Biosafety Level 2
<b>Note</b>	It is recommended to expand the cell culture and store a minimum of 10 vials at an early passage for potential future use.

---

## Materials

<b>Reagent</b>	<b>Manufacturer/Catalogue No.</b>
DMEM	Gibco/C11995500BT
Fetal Bovine Serum	ExCell/FSP500
Pen/Strep	Thermo/15140-122
G418	Genomeditech/ <a href="#">GM-040402</a>
Hygromycin	Genomeditech/ <a href="#">GM-040403</a>
Anti-CD3 hIgG1 Antibody(CH2527)	Genomeditech/ <a href="#">GM-33037AB</a>
FITC anti-human TCR α/β Antibody	Biolegend/306706
Anti-CD3 epsilon hIgG1 Antibody [OKT-3 (muromonab)]	Genomeditech/ <a href="#">GM-51478AB</a>

---

## Figures




Figure 1 | H\_CD3(TCR V2) HEK-293 Cell Line(Cat. GM-C44753) was determined by flow cytometry using Anti-CD3 hIgG1 Antibody(CH2527) (Cat. [GM-33037AB](#)).




Figure 2 | H\_CD3(TCR V2) HEK-293 Cell Line(Cat. GM-C44753) was determined by flow cytometry using FITC anti-human TCR  $\alpha/\beta$  Antibody (Biolegend/306706/B466005).




Figure 3 | H\_CD3(TCR V2) HEK-293 Cell Line(Cat. GM-C44753) was determined by flow cytometry using Anti-CD3 epsilon hlgG1 Antibody [OKT-3 (muromonab)] (Cat. [GM-51478AB](#)).

## Cell Recovery

Recovery Medium: DMEM+10% FBS+1% P.S

To insure the highest level of viability, thaw the vial and initiate the culture as soon as possible upon receipt. If upon arrival, continued storage of the frozen culture is necessary, it should be stored in liquid nitrogen vapor phase and not at -70°C. Storage at -70°C will result in loss of viability.

- Thaw the vial by gentle agitation in a 37°C water bath. To reduce the possibility of contamination, keep the O-ring and cap out of the water. Thawing should be rapid (approximately 2 - 3 minutes).
- Remove the vial from the water bath as soon as the contents are thawed, and decontaminate by dipping in or spraying with 70% ethanol. All of the operations from this point on should be carried out under strict aseptic conditions.
- Transfer the vial contents to a centrifuge tube containing 5.0 mL complete culture medium and spin at approximately 176 x g for 5 minutes. Discard supernatant.
- Resuspend cell pellet with the recommended recovery medium. And dispense into appropriate culture dishes.
- Incubate the culture at 37°C in a suitable incubator. A 5% CO<sub>2</sub> in air atmosphere is recommended if using the medium described on this product sheet.

## Cell Freezing

Freezing Medium: 90% FBS+10% DMSO

- Centrifuge at 176 x g for 3 minutes to collect cells.
- Resuspend the cells in pre-cooled freezing medium and adjust the cell density to 5E6 cells/mL.
- Aliquot 1 mL into each vial.
- Place the vial in a controlled-rate freezing container and store at -80°C for at least 1 day, then transfer to liquid nitrogen as soon as possible.

## Cell passage

Growth medium: DMEM+10% FBS+1% P.S+400 µg/mL G418+125 µg/mL Hygromycin

For the first 1 to 2 passages post-resuscitation, use the recovery medium. Once the cells have stabilized, switch to a growth medium.

- a) Subculturing is necessary when the cell density reaches 80%. It is recommended to perform subculturing at a ratio of 1:3 to 1:4 every 2-3 days. Ensure that the density does not exceed 80%, as overcrowding can lead to reduced viability due to compression.
- b) Remove and discard culture medium.
- c) Briefly rinse the cell layer with PBS to remove all traces of serum that contains trypsin inhibitor.
- d) Add 1.0 mL of 0.25% (w/v) Trypsin-EDTA solution to dish and observe cells under an inverted microscope until cell layer is dispersed (usually within 30 to 60 seconds at 37°C).
- e) Note: To avoid clumping do not agitate the cells by hitting or shaking the flask while waiting for the cells to detach. Cells that are difficult to detach may be placed at 37°C to facilitate dispersal.
- f) Add 2.0 mL of growth medium to mix well and aspirate cells by gently pipetting.
- g) After centrifugation, resuspend the pellet and add appropriate aliquots of the cell suspension to new culture vessels.
- h) Incubate cultures at 37°C.

**Subcultivation Ratio: A subcultivation ratio of 1:3 - 1:4 is recommended**

**Medium Renewal: Every 2 to 3 days**

## Notes

- a) Upon initial thawing, a higher number of dead cells is observed, which is a normal phenomenon. Significant improvement is seen after adaptation. Once the cells reach a stable state, the number of dead cells decreases after subculturing and the cell growth rate becomes stable.
- b) Ensure that the cell density does not exceed 80%, as overcrowding may lead to reduced viability due to compression.

## Related Products

CD28	
<a href="#">H_CD28 Reporter Jurkat Cell Line</a>	<a href="#">Cynomolgus_CD28 CHO-K1 Cell Line</a>
<a href="#">H_CD28 CHO-K1 Cell Line</a>	<a href="#">H_CD28 HEK-293 Cell Line</a>
<a href="#">H_CD28 KO Jurkat Cell Line</a>	
<a href="#">Anti-CD28 hIgG4 Antibody(FR104)</a>	<a href="#">Anti-H_CD28 hIgG4 Antibody(Theralizumab)</a>
<a href="#">Anti-mouse CD28 Syrian Hamster IgG2 Antibody(37. 51)</a>	
CD19	
<a href="#">H_CD19 KO Raji Cell Line</a>	<a href="#">Cynomolgus_CD19 CHO-K1 Cell Line</a>
<a href="#">Cynomolgus_CD19 HEK-293 Cell Line</a>	<a href="#">H_CD19 CHO-K1 Cell line</a>
<a href="#">H_CD19 HEK-293 Cell Line</a>	<a href="#">Mouse_CD19 CHO-K1 Cell Line</a>
<a href="#">Anti-CD19 hIgG1 Reference Antibody (Loncbio)</a>	<a href="#">Anti-CD19×CD3 hIgG1 Antibody[PIT-565(CD58 K34A)]</a>

Anti-H_CD19 hIgG1/hIgG2 Antibody(Tafasitamab)	
CD3	
ADCC FcγRIIIa(158V) Reporter Jurkat(CD3 KO) Cell Line	Jurkat CD3-BsAb Reporter Cell Line
Cynomolgus_CD3 HEK-293 Cell Line	Cynomolgus_CD3E(Membrane Bound ECD) CHO-K1 Cell Line
H_CD3 CHO-K1 Cell Line	H_CD3 HEK-293 Cell Line
H_CD3D CD3E KO Jurkat Cell Line	H_CD3E KO Jurkat Cell Line
H_CD3E(Membrane Bound ECD) CHO-K1 Cell Line	Mouse_CD3 HEK-293 Cell Line
H_CD3(TCR V2) CHO-K1 Cell Line	
Anti-CD19×CD3 hIgG1 Antibody[PIT-565(CD58 K34A)]	Anti-CD3 epsilon hIgG1 Antibody [OKT-3 (muromonab)]
Anti-CD3 hIgG1 Antibody(CH2527)	Anti-CD3×CD20 hIgG1 Bispecific Antibody (Epcobio)
Anti-CD3×FCRL5 hIgG1 Bispecific Antibody(cevostamab)	Anti-CD3E×BCMA hIgG4 Reference Antibody (Tecbio)
Anti-CD3E×DLL3 hIgG1 Bispecific Antibody(Tarlabio)	Anti-CD3E×MUC17 hIgG1 Bispecific Antibody(Vepsitbio)
Anti-mouse CD3ε mIgG2a Antibody(145-2C11)	
CD2	
H_CD2 KO Jurkat Cell Line	CD3-CD2-tsAb Reporter Jurkat(CD58 KO) Cell Line
Cynomolgus_CD2 CHO-K1 Cell Line	H_CD2 CHO-K1 Cell Line
Anti-CD19×CD3 hIgG1 Antibody[PIT-565(CD58 K34A)]	Anti-CD2 hIgG1 Antibody(BTI-322)

## License Agreement:

**By purchasing and using this cell line product, the user voluntarily agrees to accept and abide by the following policies:**

- This cell line product is restricted to research use only and shall not be used for any commercial purposes.
- This product is strictly prohibited from being used in the diagnosis or treatment of human or animal diseases, and shall not be directly used in experiments involving humans.
- Users and their contractors engaged for their benefit may use this material and its derivatives only within the agreed research scope; modification of the material is not permitted, nor may it be distributed, sold, transferred, or otherwise provided to any other entity (including affiliates).
- If use beyond the above scope is required, prior written permission from Genomeditech (Shanghai) Co.,Ltd. must be obtained. For details, please contact Genomeditech (Shanghai) Co.,Ltd.