

# Anti-Mouse\_PD1 mIgG1 Antibody

## Product Information

<b>Product Name</b>	Anti-Mouse_PD1 mIgG1 Antibody
<b>Storage temp.</b>	Store at 2-8°C short term (1-2 weeks).Store at $\leq -20^{\circ}\text{C}$ long term. Avoid repeated freeze-thaw.
<b>Catalog# / Size</b>	<b>GM-28206AB-1mg / 1 mg</b> <b>GM-28206AB-5mg / 5 mg</b> <b>GM-28206AB-25mg / 25 mg</b> <b>GM-28206AB-50mg / 50 mg</b> <b>GM-28206AB-100mg / 100 mg</b>

## Antibody Information

<b>Expression System</b>	CHO
<b>Aggregation</b>	< 5% as determined by SEC-HPLC
<b>Purity</b>	> 95% as determined by SDS-PAGE
<b>Endotoxin</b>	< 1 EU/mg, determined by LAL gel clotting assay
<b>Sterility</b>	0.2 $\mu\text{m}$ Filtered
<b>Target</b>	PD1
<b>Clone</b>	/
<b>Other Names</b>	PD-1; Pdc1; Ly101
<b>Source/Isotype</b>	Monoclonal Mouse IgG1, $\kappa$
<b>Application</b>	Flow cytometry; Binding activation: 1.5 ng/mL-100 $\mu\text{g}/\text{mL}$
<b>Description</b>	The programmed cell death 1 protein (PD-1, PDCD1, CD279) is a member of the CD28 family of immunoreceptors that regulate T cell activation and immune responses. The PD-1 protein contains an extracellular Ig V domain, a transmembrane domain, and a cytoplasmic tail that includes an immunoreceptor tyrosine-based inhibitory motif (ITIM) and an immunoreceptor tyrosine-based switch motif (ITSM). PD-1 is activated by the cell surface ligands PD-L1 and PD-L2. Upon activation, PD-1 ITIM and ITSM phosphorylation leads to the recruitment of the protein tyrosine phosphatases SHP-1 and SHP-2, which suppress TCR signaling. In addition to activated T-cells, PD-1 is expressed in activated B-cells and monocytes, although its function in these cell types has not been fully characterized. The PD-1 pathway plays an important role in immune tolerance; however, research studies show that cancer cells often adopt this pathway to escape immune surveillance. Consequently, blockade of

Version:3.1 Revision Date:15/08/2023

PD-1 and its ligands is proving to be a sound strategy for neoplastic intervention.

Formulation

Phosphate-buffered solution, pH 7.2.

## Data Examples

Flow cytometry

M\_PD1(PD-1) CHO-K1 Cell Line (Catalog # GM-C19255) was stained with Anti-Mouse\_PD1 mIgG1 Antibody (Catalog # GM-28206AB) or isotype control antibody, followed by anti-Mouse IgG FITC-conjugated Secondary Antibody.




Fig. FACS

Mouse Models





Fig 2. Mouse Tumor-Bearing Models

Binding activation

Serial dilutions of Anti-Mouse\_PD1 mIgG1 Antibody (1:4 serial dilutions, from 100 µg/mL to 1.5 ng/mL) (Catalog # GM-28206AB) were added into Mouse\_PD-1 Reporter Jurkat Cell Line (Catalog # GM-C25661). Then Mouse\_PD-1 Reporter Jurkat Cell Line was binded with Mouse PDL1 aAPC CHO-K1 Cell Line (Catalog # GM-C25791).EC<sub>50</sub> for this effect was 4.126 µg/mL.



M_PD-1 PDL1 Reporter Blockade Assay	
EC50	4.126

Fig 3. assay