Toll-free: 400 627 9288

Email: service@genomeditech.com



Anti-Mouse_PD1 mlgG1 Antibody(29F.1A12)

Product Information

Product Name Anti-Mouse_PD1 mlgG1 Antibody(29F.1A12)

Storage temp. Store at 2-8°C short term (1-2 weeks). Store at \leq -20°C long term. Avoid

repeated freeze-thaw.

Catalog# / Size GM-86949MAB-1mg / 1 mg

GM-86949MAB-5mg / 5 mg GM-86949MAB-25mg / 25 mg GM-86949MAB-50mg / 50 mg GM-86949MAB-100mg / 100 mg

Antibody Information

Expression System CHO

Aggregation < 5% as determined by SEC-HPLC

Purity > 95% as determined by SDS-PAGE

Endotoxin < 1 EU/mg, determined by LAL gel clotting assay

Sterility 0.2 µm Filtered

 Target
 PD1

 Clone
 29F.1A12

Other Names PD-1; Pdc1; Ly101

Source/Isotype Monoclonal mouse IgG1 D265A, kappa

Application /

Description The programmed cell death 1 protein (PD-1, PDCD1, CD279) is a member

of the CD28 family of immunoreceptors that regulate T cell activation and immune responses. The PD-1 protein contains an extracellular Ig V domain, a transmembrane domain, and a cytoplasmic tail that includes an immunoreceptor tyrosine-based inhibitory motif (ITIM) and an immunoreceptor tyrosine-based switch motif (ITSM). PD-1 is activated by the cell surface ligands PD-L1 and PD-L2. Upon activation, PD-1 ITIM and ITSM phosphorylation leads to the recruitment of the protein tyrosine phosphatases SHP-1 and SHP-2, which suppress TCR signaling. In addition to activated T-cells, PD-1 is expressed in activated B-cells and monocytes, although its function in these cell types has not been fully characterized. The PD-1 pathway plays an important role in immune tolerance; however, research studies show that cancer cells often adopt this pathway to escape immune surveillance. Consequently, blockade of

Version:3.1

Genomeditech (Shanghai) Co.,Ltd. Order: 021-68455258/50432826/50432825

Toll-free: 400 627 9288

Email: service@genomeditech.com

PD-1 and its ligands is proving to be a sound strategy for neoplastic


intervention.

Formulation

Phosphate-buffered solution, pH 7.2.


Data Examples

SDS-PAGE



On SDS-PAGE under reducing (R)/non-reducing (N-R) condition. The gel was stained overnight with Coomassie Blue. The purity of the protein is greater than 95%.

SEC-HPLC



The purity of this product is more than 95% verified by SEC-HPLC.