

Anti-NPR1 hlgG4 Antibody(REGN-5381)

Product information

GM-87697AB-10	10 µg
GM-87697AB-100	100 µg
GM-87697AB-1000	1 mg

Antibody Information


Species Reactivity	Human;
Clone	REGN-5381
Source/Isotype	Monoclonal human IgG4, κ
Application	Flow cytometry
Specificity	Detects NPR1
Gene	NPR1
Other Names	ANPRA, ANPa, GUC2A, GUCY2A, NPRA
Gene ID	4881(human);
Background	Guanylyl cyclases, catalyzing the production of cGMP from GTP, are classified as soluble and membrane forms. The membrane guanylyl cyclases, often termed guanylyl cyclases A through F, form a family of cell-surface receptors with a similar topographic structure: an extracellular ligand-binding domain, a single membrane-spanning domain, and an intracellular region that contains a protein kinase-like domain and a cyclase catalytic domain. GC-A and GC-B function as receptors for natriuretic peptides; they are also referred to as atrial natriuretic peptide receptor A (NPR1) and type B (NPR2).which encodes a protein with only the ligand-binding transmembrane and 37-amino acid cytoplasmic domains. NPR1 is a membrane-bound guanylate cyclase that serves as the receptor for both atrial and brain natriuretic peptides (ANP) and BNP, respectively).
Storage	Store at 2-8°C short term (1-2 weeks).Store at ≤ -20°C long term. Avoid repeated freeze-thaw.
Formulation	Phosphate-buffered solution, pH 7.2.
Endotoxin	< 1 EU/mg, determined by LAL gel clotting assay

Version:3.2 Revision Date:03/25/2024

Data Examples

Flow cytometry

H_NPR1 HEK-293 Cell Line(Catalog # GM-C34844) was stained with Anti-NPR1 hIgG4 Antibody(REGN-5381) (Catalog # GM-87697AB) or isotype control antibody, followed by anti-Human IgG APC-conjugated Secondary Antibody.



SampleID	Geometric Mean : FL11-H
HEK-293 anti-NPR1+APC-2nd	4387
HEK-293 H_NPR1 H_IgG+APC-2nd Ab	1899
HEK-293 H_NPR1 anti-NPR1+APC-2nd Ab	6.94E5

Fig. FACS