

Product Sheet


H_FOXP3-Promoter Reporter Jurkat Cell Line

Catalog number: GM-C13108

Version 3.3.1.241226

FOXP3 belongs to the forkhead family and is a major transcription factor. Studies have shown that overexpressed FOXP3 in Jurkat cells can compete with NFAT for binding or directly interact with NFAT or NF-κB as a transcriptional repressor of the IL-2 promoter. In humans, the expression of FOXP3 is upregulated in all activated T cells. FOXP3 is essential for the development and function of naturally occurring Treg cells and is involved in the molecular mechanisms that control immune tolerance.

H_FOXP3-Promoter Reporter Jurkat Cell Line is a clonal stable cell line constructed using lentiviral technology that expresses a FOXP3-Promoter inducible luciferase reporter gene. When PMA and Ionomycin activates downstream signaling pathways, it leads to the expression of luciferase driven by the FOXP3-Promoter. The luciferase activity measurement indicates the activation level of the signaling pathway and can thus be used to evaluate the in vitro effects of drugs related to FOXP3-Promoter.



Specifications

Quantity	5E6 Cells per vial,1 mL
Product Format	1 vial of frozen cells
Shipping	Shipped on dry ice
Storage Conditions	Liquid nitrogen immediately upon receipt

Recovery Medium	RPMI 1640+10% FBS+1% P.S
Growth medium	RPMI 1640+10% FBS+1% P.S+0.75 µg/mL Puromycin
Note	None
Freezing Medium	90% FBS+10% DMSO
Growth properties	Suspension
Growth Conditions	37°C, 5% CO ₂

Mycoplasma Testing	The cell line has been screened to confirm the absence of Mycoplasma species.
Safety considerations	Biosafety Level 2
Note	It is recommended to expand the cell culture and store a minimum of 10 vials at an early passage for potential future use.

Materials

Reagent	Manufacturer/Catalogue No.
RPMI 1640	gibco/C11875500BT
Fetal Bovine Serum	Cegrogen biotech/A0500-3010
Pen/Strep	Thermo/15140-122
Puromycin	Genomeditech/ GM-040401
PMA/TPA	Beyotime/S1819
Ionomycin	MCE/HY-13434
GMOne-Step Luciferase Reporter Gene Assay Kit	Genomeditech/ GM-040503

Figures




Figure 1 | Response to PMA/TPA+Ionomycin. The H_FOXP3-Promoter Reporter Jurkat Cell Line (Cat. GM-C13108) at a concentration of 1E5 cells/well (96-well format) was stimulated with serial dilutions of PMA/TPA (Beyotime/S1819) and 0.5 µg/mL Ionomycin (MCE/HY-13434) in assay buffer (RPMI 1640 + 1% FBS + 1% P.S) for 7 hours. The firefly luciferase activity was measured using the GMOne-Step Luciferase Reporter Gene Assay Kit (Cat. GM-040503). The maximum induction fold was approximately [16.9]. Data are shown by drug mass concentration.




Figure 2 | Response to PMA/TPA+Ionomycin. The H_FOXP3-Promoter Reporter Jurkat Cell Line (Cat. GM-C13108) at a concentration of 1E5 cells/well (96-well format) was stimulated with serial dilutions of PMA/TPA (Beyotime/S1819) and 0.5 µg/mL Ionomycin (MCE/HY-13434) in assay buffer (RPMI 1640 + 1% FBS + 1% P.S) for 7 hours. The firefly luciferase activity was measured using the GMOne-Step Luciferase Reporter Gene Assay Kit (Cat. GM-040503). The maximum induction fold was approximately [16.9]. Data are shown by drug molar concentration.

Cell Recovery

Recovery Medium: RPMI 1640+10% FBS+1% P.S

To insure the highest level of viability, thaw the vial and initiate the culture as soon as possible upon receipt. If upon arrival, continued storage of the frozen culture is necessary, it should be stored in liquid nitrogen vapor phase and not at -70°C . Storage at -70°C will result in loss of viability.

- Thaw the vial by gentle agitation in a 37°C water bath. To reduce the possibility of contamination, keep the O-ring and cap out of the water. Thawing should be rapid (approximately 2 - 3 minutes).
- Remove the vial from the water bath as soon as the contents are thawed, and decontaminate by dipping in or spraying with 70% ethanol. All of the operations from this point on should be carried out under strict aseptic conditions.
- Transfer the vial contents to a centrifuge tube containing 5.0 mL complete culture medium. And spin at approximately $176 \times g$ for 5 minutes. Discard supernatant.
- Resuspend cell pellet with the recommended complete medium. And dispense the suspension into 1 - 2 T-25 culture flasks.
- Incubate the culture at 37°C in a suitable incubator. A 5% CO_2 in air atmosphere is recommended if using the medium described on this product sheet.

Cell Freezing

Freezing Medium: 90% FBS+10% DMSO

- Centrifuge at $176 \times g$ for 3 minutes to collect cells.
- Resuspend the cells in pre-cooled freezing medium and adjust the cell density to 5×10^6 cells/mL.
- Aliquot 1 mL into each vial.
- Place the vial in a controlled-rate freezing container and store at -80°C for at least 1 day, then transfer to liquid nitrogen as soon as possible.

Cell passage

Growth medium: RPMI 1640+10% FBS+1% P.S+0.75 $\mu\text{g}/\text{mL}$ Puromycin

Approximately 48-72 hours after the initial thawing, the cells can be passaged for the first time. After this initial passage, the culture medium can be adjusted to growth medium supplemented with antibiotics. If cells are not passaged within 48 hours, it is recommended to add some fresh recovery medium and place the flask horizontally.

- When the cell density reaches $1.5 - 2 \times 10^6$ cells/mL, subculture the cells. Do not allow the cell density to exceed 2×10^6 cells/mL.
- It is recommended to use T-25 flasks for subculturing.
- These cells are suspension cells, and it is recommended to use the "half-medium change" method to maintain optimal cell conditions during passaging.
- During passaging, you can directly add fresh growth medium to the culture flask, gently pipette to resuspend the cells, and then transfer the cell suspension to a new T-25 flask for continued culture.

Subcultivation Ratio: Maintain cultures at a cell concentraion between 3×10^5 and 1×10^6 viable cells/mL.

Medium Renewal: Every 2 to 3 days

Notes

- These cells are sensitive to density, so please ensure that the cell density is maintained within an appropriate range during culture and subculturing.
- During the first passage, pay attention to the nutrient supply; if not subculturing, make sure to add fresh recovery medium every other day as needed.

Related Products

TCR	
H_IL2-Promoter Reporter Jurkat Cell Line	NFAT-Luc Reporter Jurkat Cell Line
TCR Knockout Reporter Cell Line(CD4+)	OKT3(CD3 ScFv) CHO-K1 Cell Line
Anti-CD3-CD19 Bispecific Antibody(Blinatumomab)	

Limited Use License Agreement

Genomeditech (Shanghai) Co., Ltd grants to the Licensee all intellectual property rights, exclusive, non-transferable, and non-sublicensable rights of the Licensed Materials; Genomeditech (Shanghai) Co., Ltd will retain ownership of the Licensed Materials, cell line history packages, progeny, and the Licensed Materials including modified materials.

Between Genomeditech (Shanghai) Co., Ltd, and Licensee, Licensee is not permitted to modify cell lines in any way. The Licensee shall not share, distribute, sell, sublicense, or otherwise provide the Licensed Materials, or progenitors to third parties such as laboratories, departments, research institutions, hospitals, universities, or biotechnology companies for use other than for the purpose of outsourcing the Licensee's research.

Please refer to the Genomeditech Cell Line License Agreement for details.