

# Product Sheet

## H\_HLA-G1 PDL1 MC38(mouse\_PDL1 KO) Cell Line

Catalog number: GM-C22142

Version 3.3.1.250126

<b>Description</b>	H_HLA-G1 PDL1 MC38(mouse_PDL1 KO) Cell Line is a clonal stable MC38 cell line that continuously expresses human HLA-G1, human PDL1 and human B2M. It is constructed using lentiviral technology, based on the knockout of mouse PDL1.
<b>Quantity</b>	5E6 Cells per vial, 1 mL
<b>Product Format</b>	3 vials of frozen cells
<b>Shipping</b>	Shipped on dry ice
<b>Storage Conditions</b>	Liquid nitrogen immediately upon receipt
<b>Target</b>	Human_HLA-G1 & Human_PDL1 & Human_B2M
<b>Gene ID/Uniprot ID</b>	NP_002118.1 & NP_054862.1 & P61769
<b>Host Cell</b>	MC38
<b>Recovery Medium</b>	DMEM+10% FBS+1% P.S
<b>Growth medium</b>	DMEM+10% FBS+1% P.S+2 µg/mL Blasticidin+500 µg/mL Bleomycin+200 µg/mL G418+200 µg/mL Hygromycin+2.5 µg/mL Puromycin
<b>Note</b>	None
<b>Freezing Medium</b>	90% FBS+10% DMSO
<b>Growth properties</b>	Adherent
<b>Growth Conditions</b>	37°C, 5% CO <sub>2</sub>
<b>Mycoplasma Testing</b>	The cell line has been screened to confirm the absence of Mycoplasma species.
<b>Safety considerations</b>	Biosafety Level 2
<b>Note</b>	It is recommended to expand the cell culture and store a minimum of 10 vials at an early passage for potential future use.

## Materials

Reagent	Manufacturer/Catalogue No.
DMEM	VivaCell/C3110-0500
Fetal Bovine Serum	Cegrogen biotech/A0500-3010
Pen/Strep	Thermo/15140-122
Blasticidin	Genomeditech/GM-040404
Bleomycin	Genomeditech/GM-040407
G418	Genomeditech/GM-040402
Hygromycin	Genomeditech/GM-040403
Puromycin	Genomeditech/GM-040401
Anti-H_HLA-G1 hIgG1 Antibody(38373)	Genomeditech/GM-28208AB
Anti-H_CD274(PDL1) hIgG1 Antibody(Atezolizumab)	Genomeditech/GM-31740AB

## Figures




Figure 1 | H\_HLA-G1 PDL1 MC38(mouse\_PDL1 KO) Cell Line (Cat. GM-C22142) was determined by flow cytometry using Anti-H\_HLA-G1 hIgG1 Antibody(38373) (Cat. [GM-28208AB](#)).




Figure 2 | H\_HLA-G1 PDL1 MC38(mouse\_PDL1 KO) Cell Line (Cat. GM-C22142) was determined by flow cytometry using Anti-H\_CD274(PDL1) hIgG1 Antibody(Atezolizumab) (Cat. GM-31740AB).




Figure 3 | H\_HLA-G1 PDL1 MC38(mouse\_PDL1 KO) Cell Line (Cat. GM-C22142) was determined by flow cytometry using PE anti-mouse CD274 (B7-H1, PD-L1) Antibody (BioLegend/124307).




Figure 4 | The Sanger sequencing of the H\_HLA-G1 PDL1 MC38(mouse\_PDL1 KO) Cell Line (Cat. GM-C22142) showed successful knockout of PDL1.

## Cell Recovery

Recovery Medium: DMEM+10% FBS+1% P.S

To insure the highest level of viability, thaw the vial and initiate the culture as soon as possible upon receipt. If upon arrival, continued storage of the frozen culture is necessary, it should be stored in liquid nitrogen vapor phase and not at  $-70^{\circ}\text{C}$ . Storage at  $-70^{\circ}\text{C}$  will result in loss of viability.

- a) Thaw the vial by gentle agitation in a  $37^{\circ}\text{C}$  water bath. To reduce the possibility of contamination, keep the O-ring and cap out of the water. Thawing should be rapid (approximately 2 - 3 minutes).
- b) Remove the vial from the water bath as soon as the contents are thawed, and decontaminate by dipping in or spraying with 70% ethanol. All of the operations from this point on should be carried out under strict aseptic conditions.
- c) Transfer the vial contents to a centrifuge tube containing 5.0 mL complete culture medium and spin at approximately  $176 \times g$  for 5 minutes. Discard supernatant.
- d) Resuspend cell pellet with the recommended recovery medium. And dispense into appropriate culture dishes.
- e) Incubate the culture at  $37^{\circ}\text{C}$  in a suitable incubator. A 5%  $\text{CO}_2$  in air atmosphere is recommended if using the medium described on this product sheet.

## Cell Freezing

Freezing Medium: 90% FBS+10% DMSO

- a) Centrifuge at  $176 \times g$  for 3 minutes to collect cells.
- b) Resuspend the cells in pre-cooled freezing medium and adjust the cell density to  $5 \times 10^6$  cells/mL.
- c) Aliquot 1 mL into each vial.
- d) Place the vial in a controlled-rate freezing container and store at  $-80^{\circ}\text{C}$  for at least 1 day, then transfer to liquid nitrogen as soon as possible.

## Cell passage

Growth medium: DMEM+10% FBS+1% P.S+2  $\mu\text{g}/\text{mL}$  Blasticidin+500  $\mu\text{g}/\text{mL}$  Bleomycin+200  $\mu\text{g}/\text{mL}$  G418+200  $\mu\text{g}/\text{mL}$  Hygromycin+2.5  $\mu\text{g}/\text{mL}$  Puromycin

For the first 1 to 2 passages post-resuscitation, use the recovery medium. Once the cells have stabilized, switch to a growth medium.

- a) Remove and discard culture medium.
- b) Briefly rinse the cell layer with PBS to remove all traces of serum that contains trypsin inhibitor.
- c) Add 1.0 mL of 0.25% (w/v) Trypsin-EDTA solution to dish and observe cells under an inverted microscope until cell layer is dispersed (usually within 30 to 60 seconds at  $37^{\circ}\text{C}$ ).
- d) Note: To avoid clumping do not agitate the cells by hitting or shaking the flask while waiting for the cells to detach. Cells that are difficult to detach may be placed at  $37^{\circ}\text{C}$  to facilitate dispersal.
- e) Add 2.0 mL of growth medium to mix well and aspirate cells by gently pipetting.

- f) After centrifugation, resuspend the pellet and add appropriate aliquots of the cell suspension to new culture vessels.  
g) Incubate cultures at 37°C.

**Subcultivation Ratio: A subcultivation ratio of 1:4 - 1:5 is recommended**

**Medium Renewal: Every 2 to 3 days**

## Notes

- a) After the stabilization of the cell condition, there will be fewer dead cells post-passage, the cell growth rate will tend to stabilize, cell morphology will become uniform, and the cells will appear robust.

## Sequence

### HLA-G1 NP\_002118.1

MVVMAPRTLFLLLSGALTLTETWAGSHSMRYFSAAVSRPGRGEPFIAMGYVDDTQFVRFSDSACPRMEP  
RAPWVEQEGPEYWEEETRNTKAHAQTDRMNLQTLRGYYNQSEASSHTLQWMIGCDLGS DGRLLRGYEQY  
AYDGKDY LALNEDLRSWTAADTAAQISKRKCEANVAEQRRAYLEGTCVEWLHRYLENGKEMLRADPP  
KTHVTHHPVFDYEATLRCWALGFYPAEIIITWQRDGEDQTQDVELVETRPAGDGTQKWA AVVVPSGEEQR  
YTCHVQHEGLPEPLMLRWKQSSSLPTIPIMGIVAGLVVLA AVVTGAAVA AVLWRKKSSD\*

### CD274(PD-L1) NP\_054862.1

MRIFAVFIFMTYWHLNNAFTVTVPKDLVVEYGSNMTECKFPVEKQLDLAALIVYWEMEDKNIIQFVHGEE  
DLKVQHSSYRQRARLLKDQLSLGNAALQITDVKLQDAGVYRCMISYGGADYKRITVKVNAPYNKINQRILV  
VDPVTSEHELTCQAEGYPKAEVIWTSSDHQVLSGKTTTTNSKREEKLFNVTSTLRINTTTNEIFYCTFRRLDPE  
ENHTAELVIPPLAHPPNERHLVILGAILLCLGVALTFIFRLRKGRMMDVKKCGIQDTNSKKQSDTHLEET\*

### B2M P61769

MSRSVALAVLALLSLSGLEAIQRTPKIQVYSRHPAENGKSNFLNCYVSGFHPSDIEVDLLKNGERIEKVEHSDL  
SFSKDWSFYLLYYTEFTPTEKDEYACRVNHVTLSPKIVKWDRDM\*

## Related Products

LILRB1(ILT2)	
<a href="#">H_LILRB1(ILT2) Reporter Jurkat Cell Line</a>	<a href="#">H_LILRB1(ILT2) CHO-K1 Cell Line</a>
<a href="#">H_LILRB1(ILT2) HEK-293 Cell Line</a>	<a href="#">Rhesus_LILRB1 CHO-K1 Cell Line</a>
<a href="#">Anti-LILRB1(ILT2) mIgG1 Antibody(12D12)</a>	
HLA-G	
<a href="#">H_HLA-G1 CHO-K1 Cell Line</a>	<a href="#">H_HLA-G1 HEK-293 Cell Line</a>
<a href="#">H_HLA-G1 MC38 Cell Line</a>	<a href="#">H_HLA-G1 OKT3 CHO-K1 Cell Line</a>
<a href="#">H_HLA-G1 SK-OV3 Cell Line</a>	<a href="#">Rhesus_MAMU-AG HEK-293 Cell Line</a>
<a href="#">Anti-H_HLA-G1 hIgG1 Antibody(38373)</a>	
In Vivo MAb Isotype Controls	
<a href="#">Human IgG1 Isotype Control(Anti-HEL)</a>	<a href="#">Human IgG1 Isotype Control(Anti-MOPC-21)</a>

Human IgG1 Isotype Control(Anti-RSV)	Human IgG1(LALA) Isotype Control(Anti-HEL)
Human IgG1(LALAPG) Isotype Control(Anti-HEL)	Human IgG1(N297A) Isotype Control(Anti-HEL)
Human IgG4(S228P) Isotype Control(Anti-HEL)	Mouse IgG1 Isotype Control(Anti-HEL)
Mouse IgG2a Isotype Control(Anti-HEL)	Mouse IgG2a Isotype Control(Anti-RSV)
Mouse IgG2a(D265A) Isotype Control(Anti-HEL)	

## Limited Use License Agreement

Genomeditech (Shanghai) Co., Ltd grants to the Licensee all intellectual property rights, exclusive, non-transferable, and non-sublicensable rights of the Licensed Materials; Genomeditech (Shanghai) Co., Ltd will retain ownership of the Licensed Materials, cell line history packages, progeny, and the Licensed Materials including modified materials.

Between Genomeditech (Shanghai) Co., Ltd, and Licensee, Licensee is not permitted to modify cell lines in any way. The Licensee shall not share, distribute, sell, sublicense, or otherwise provide the Licensed Materials, or progenitors to third parties such as laboratories, departments, research institutions, hospitals, universities, or biotechnology companies for use other than for the purpose of outsourcing the Licensee's research.

Please refer to the Genomeditech Cell Line License Agreement for details.

Genomeditech