

Product Sheet

ADCC FcγRIIIa(158V) DDX35™ Jurkat Effector Cell Line


Catalog number: GM-C26024

Version 3.3.1.241212

ADCC, or antibody-dependent cell-mediated cytotoxicity, is a process where immune cells with Fc receptors kill target cells that bind to antibodies via the Fc region. This mechanism is used to assess the efficacy of antibodies and target cells. Antibodies attach to antigens on target cell surfaces, and when the Fc region binds to the FcγRIIIa receptor on effector cells (mainly natural killer cells), it triggers cross-linking and activates the ADCC signaling pathway. The 158V variant is a polymorphism where valine (V) replaces phenylalanine (F) at position 158, resulting in higher affinity for antibodies.

ADCC FcγRIIIa(158V) DDX35™ Jurkat Effector Cell Line is a clonal stable Jurkat cell line constructed using lentiviral technology, constitutive expression of the FcγRIIIa(158V) gene, along with signal-dependent expression of a luciferase reporter gene. When IgG binds to target cells and effector cells, it leads to the expression of luciferase, which can be used to evaluate the biological activity of antibodies in the mechanism of ADCC.

ADCC FcγRIIIa(158V) DDX35™ Jurkat Effector Cell Line was obtained through extensive monoclonal screening and multiple rounds of monoclonal selection. It possesses high stability, high sensitivity, and high amplification properties, meeting the standards for customers' batch library construction and release experiments.



Specifications

Quantity	5E6 Cells per vial,1 mL
Product Format	1 vial of frozen cells
Shipping	Shipped on dry ice
Storage Conditions	Liquid nitrogen immediately upon receipt

Recovery Medium	RPMI 1640+10% FBS+1% P.S
Growth medium	RPMI 1640+10% FBS+1% P.S+3.5 µg/mL Blasticidin+0.75 µg/mL Puromycin
Note	None
Freezing Medium	90% FBS+10% DMSO
Growth properties	Suspension
Growth Conditions	37°C, 5% CO ₂

Mycoplasma Testing	The cell line has been screened to confirm the absence of Mycoplasma species.
Safety considerations	Biosafety Level 2
Note	It is recommended to expand the cell culture and store a minimum of 10 vials at an early passage for potential future use.

Materials

Reagent	Manufacturer/Catalogue No.
RPMI 1640	VivaCell/C3010-0500
Fetal Bovine Serum	Cegrogen biotech/A0500-3010
Pen/Strep	Thermo/15140-122
Blasticidin	Genomeditech/ GM-040404
Puromycin	Genomeditech/ GM-040401
H_CLDN18.2(isoform2) CHO-K1 Cell Line	Genomeditech/ GM-C05273
Anti-CLDN18.2 hIgG1 Antibody(Zolbetuximab)	Genomeditech/ GM-34137AB
GMOne-Step Luciferase Reporter Gene Assay Kit	Genomeditech/ GM-040503

Figures




Figure 1 | The passage stability of response to Anti-CLDN18.2 hIgG1 Antibody. Serial dilutions of the Anti-CLDN18.2 hIgG1 Antibody(Zolbetuximab) (Cat. [GM-34137AB](#)) and 1.5E5 cells/well of the passage 5, 10, 20, 25, 30 and 35 of the ADCC FcγRIIIa(158V) DDX35™ Jurkat Effector Cell Line (Cat. [GM-C26024](#)) were added to 1E4 cells/well of H_CLDN18.2 CHO-K1 cell line (Cat. [GM-C05273](#)) for 24 hours. Firefly luciferase activity is then measured using the GMOne-Step Luciferase Reporter Gene Assay Kit (Cat. [GM-040503](#)). Data are shown by drug mass concentration.




Figure 2 | The passage stability of response to Anti-CLDN18.2 hIgG1 Antibody. Serial dilutions of the Anti-CLDN18.2 hIgG1 Antibody(Zolbetuximab) (Cat. [GM-34137AB](#)) and 1.5E5 cells/well of the passage 8 and 38 of the ADCC FcγRIIIa(158V) DDX35™ Jurkat Effector Cell Line (Cat. [GM-C26024](#)) were added to 1E4 cells/well of H_CLDN18.2 CHO-K1 cell line (Cat. [GM-C05273](#)) for 24 hours. Firefly luciferase activity is then measured using the GMOne-Step Luciferase Reporter Gene Assay Kit (Cat. [GM-040503](#)). Data are shown by drug mass concentration.




Figure 3 | Response to Anti-CLDN18.2 hIgG1 Antibody. Serial dilutions of the Anti-CLDN18.2 hIgG1 Antibody(Zolbetuximab) (Cat. [GM-34137AB](#)) and 1.5E5 cells/well of the ADCC FcγRIIIa(158V) DDX35™ Jurkat Effector Cell Line (Cat. GM-C26024) were added to 1E4 cells/well of H_CLDN18.2 CHO-K1 cell line (Cat. [GM-C05273](#)) for 24 hours. Firefly luciferase activity is then measured using the GMOne-Step Luciferase Reporter Gene Assay Kit (Cat. [GM-040503](#)). The maximum induction fold was approximately[16.6]. Data are shown by drug mass concentration.




Figure 4 | Response to Anti-CLDN18.2 hIgG1 Antibody. Serial dilutions of the Anti-CLDN18.2 hIgG1 Antibody(Zolbetuximab) (Cat. [GM-34137AB](#)) and 1.5E5 cells/well of the ADCC FcγRIIIa(158V) DDX35™ Jurkat Effector Cell Line (Cat. GM-C26024) were added to 1E4 cells/well of H_CLDN18.2 CHO-K1 cell line (Cat. [GM-C05273](#)) for 24 hours. Firefly luciferase activity is then measured using the GMOne-Step Luciferase Reporter Gene Assay Kit (Cat. [GM-040503](#)). The maximum induction fold was approximately[16.6]. Data are shown by drug molar concentration.

Cell Recovery

Recovery Medium: RPMI 1640+10% FBS+1% P.S

To insure the highest level of viability, thaw the vial and initiate the culture as soon as possible upon receipt. If upon arrival, continued storage of the frozen culture is necessary, it should be stored in liquid nitrogen vapor phase and not at -70°C . Storage at -70°C will result in loss of viability.

- a) Thaw the vial by gentle agitation in a 37°C water bath. To reduce the possibility of contamination, keep the O-ring and cap out of the water. Thawing should be rapid (approximately 2 - 3 minutes).
- b) Remove the vial from the water bath as soon as the contents are thawed, and decontaminate by dipping in or spraying with 70% ethanol. All of the operations from this point on should be carried out under strict aseptic conditions.
- c) Transfer the vial contents to a centrifuge tube containing 5.0 mL complete culture medium. And spin at approximately $176 \times g$ for 5 minutes. Discard supernatant.
- d) Resuspend cell pellet with the recommended complete medium. And dispense the suspension into 1 - 2 T-25 culture flasks.
- e) Incubate the culture at 37°C in a suitable incubator. A 5% CO_2 in air atmosphere is recommended if using the medium described on this product sheet.

Cell Freezing

Freezing Medium: 90% FBS+10% DMSO

- a) Centrifuge at $176 \times g$ for 3 minutes to collect cells.
- b) Resuspend the cells in pre-cooled freezing medium and adjust the cell density to 5×10^6 cells/mL.
- c) Aliquot 1 mL into each vial.
- d) Place the vial in a controlled-rate freezing container and store at -80°C for at least 1 day, then transfer to liquid nitrogen as soon as possible.

Cell passage

Growth medium: RPMI 1640+10% FBS+1% P.S+3.5 $\mu\text{g}/\text{mL}$ Blasticidin+0.75 $\mu\text{g}/\text{mL}$ Puromycin

Approximately 48-72 hours after the initial thawing, the cells can be passaged for the first time. After this initial passage, the culture medium can be adjusted to growth medium supplemented with antibiotics. If cells are not passaged within 48 hours, it is recommended to add some fresh recovery medium and place the flask horizontally.

- a) When the cell density reaches $1.5 - 2 \times 10^6$ cells/mL, subculture the cells. Do not allow the cell density to exceed 2×10^6 cells/mL.
- b) It is recommended to use T-25 flasks for subculturing.
- c) These cells are suspension cells, and it is recommended to use the "half-medium change" method to maintain optimal cell conditions during passaging.
- d) During passaging, you can directly add fresh growth medium to the culture flask, gently pipette to resuspend the cells, and then transfer the cell suspension to a new T-25 flask for continued culture.

Subcultivation Ratio: Maintain cultures at a cell concentraion between 3E5 and 1E6 viable cells/mL.

Medium Renewal: Every 2 to 3 days

Notes

- These cells are sensitive to density, so please ensure that the cell density is maintained within an appropriate range during culture and subculturing.
- During the first passage, pay attention to the nutrient supply; if not subculturing, make sure to add fresh recovery medium every other day as needed.

Related Products

FcγR	
Cynomolgus_FcRn MDCK Cell Line	H_FCGR1A(CD64) CHO-K1 Cell Line
H_FCGR1A(CD64) HEK-293 Cell Line	H_FCGR2A(CD32A) CHO-K1 Cell Line
H_FCGR2B(CD32B) CHO-K1 Cell Line	H_FCGR3A(CD16a) 158F CHO-K1 Cell Line
H_FCGR3A(CD16a) 158V CHO-K1 Cell Line	H_FCGR3B(CD16b) CHO-K1 Cell Line
H_FcRn CHO-K1 Cell Line	H_FcRn MDCK Cell Line
Mouse_FcRn MDCK Cell Line	
Anti-FcRn hIgG4 Reference Antibody(Rozabio)	Anti-H_FcRn IgG4 Antibody(Rozanolixizumab)
Anti-Mouse CD1632 mIgG2b Antibody(2.4G2)	
ADCCP	
ADCC FcγRIIIa(158F) Jurkat Effector Cell Line	ADCC FcγRIIIa(158V) Jurkat Effector Cell Line
ADCC M_FcγRIV Jurkat Effector Cell Line	ADCP FcγRIIa DDX35TM Jurkat Effector Cell Line
ADCP FcγRIIa Jurkat Effector Cell Line	ADCP FcγRIIa R131 Jurkat Effector Cell Line
ADCP FcγRIIb Jurkat Effector Cell Line	

Limited Use License Agreement

Genomeditech (Shanghai) Co., Ltd grants to the Licensee all intellectual property rights, exclusive, non-transferable, and non-sublicensable rights of the Licensed Materials; Genomeditech (Shanghai) Co., Ltd will retain ownership of the Licensed Materials, cell line history packages, progeny, and the Licensed Materials including modified materials.

Between Genomeditech (Shanghai) Co., Ltd, and Licensee, Licensee is not permitted to modify cell lines in any way. The Licensee shall not share, distribute, sell, sublicense, or otherwise provide the Licensed Materials, or progenitors to third parties such as laboratories, departments, research institutions, hospitals, universities, or biotechnology companies for use other than for the purpose of outsourcing the Licensee's research.

Please refer to the Genomeditech Cell Line License Agreement for details.