

# Product Sheet

## H\_CDH17 RKO Cell Line

Catalog number: GM-C31740

Version 3.3.1.241219

<b>Description</b>	H_CDH17 RKO Cell Line is a clonal stable RKO cell line that constitutively expresses the human CDH17 gene, constructed using lentiviral technology.
<b>Quantity</b>	5E6 Cells per vial, 1 mL
<b>Product Format</b>	1 vial of frozen cells
<b>Shipping</b>	Shipped on dry ice
<b>Storage Conditions</b>	Liquid nitrogen immediately upon receipt
<b>Target</b>	Human_CDH17
<b>Gene ID/Uniprot ID</b>	Q12864(AA Met 1 - Ile 808)
<b>Host Cell</b>	RKO
<b>Recovery Medium</b>	DMEM+10% FBS+1% P.S
<b>Growth medium</b>	DMEM+10% FBS+1% P.S+15 µg/mL Blasticidin+0.25 µg/mL Puromycin
<b>Note</b>	None
<b>Freezing Medium</b>	90% FBS+10% DMSO
<b>Growth properties</b>	Adherent
<b>Growth Conditions</b>	37°C, 5% CO <sub>2</sub>
<b>Mycoplasma Testing</b>	The cell line has been screened to confirm the absence of Mycoplasma species.
<b>Safety considerations</b>	Biosafety Level 2
<b>Note</b>	It is recommended to expand the cell culture and store a minimum of 10 vials at an early passage for potential future use.

## Materials

Reagent	Manufacturer/Catalogue No.
DMEM	Gibco/C11995500BT
Fetal Bovine Serum	Cegrogen biotech/A0500-3010
Pen/Strep	Thermo/15140-122
Blasticidin	Genomeditech/GM-040404
Puromycin	Genomeditech/GM-040401
Anti-CDH17 hIgG1 Antibody(BI-905711)	Genomeditech/GM-52672AB

## Figures




Figure 1 | H\_CDH17 RKO Cell Line (Cat. GM-C31740) was determined by flow cytometry using Anti-CDH17 hIgG1 Antibody(BI-905711) (Cat. [GM-52672AB](#)).




Figure 2 | The passage stability of the H\_CDH17 RKO Cell Line (Cat. GM-C31740) was determined by flow cytometry using Anti-CDH17 hIgG1 Antibody(BI-905711) (Cat. [GM-52672AB](#)).

## Cell Recovery

Recovery Medium: DMEM+10% FBS+1% P.S

To insure the highest level of viability, thaw the vial and initiate the culture as soon as possible upon receipt. If upon arrival, continued storage of the frozen culture is necessary, it should be stored in liquid nitrogen vapor phase and not at  $-70^{\circ}\text{C}$ . Storage at  $-70^{\circ}\text{C}$  will result in loss of viability.

- Thaw the vial by gentle agitation in a  $37^{\circ}\text{C}$  water bath. To reduce the possibility of contamination, keep the O-ring and cap out of the water. Thawing should be rapid (approximately 2 - 3 minutes).
- Remove the vial from the water bath as soon as the contents are thawed, and decontaminate by dipping in or spraying with 70% ethanol. All of the operations from this point on should be carried out under strict aseptic conditions.
- Transfer the vial contents to a centrifuge tube containing 5.0 mL complete culture medium and spin at approximately  $176 \times g$  for 5 minutes. Discard supernatant.
- Resuspend cell pellet with the recommended recovery medium. And dispense into appropriate culture dishes.
- Incubate the culture at  $37^{\circ}\text{C}$  in a suitable incubator. A 5%  $\text{CO}_2$  in air atmosphere is recommended if using the medium described on this product sheet.

## Cell Freezing

Freezing Medium: 90% FBS+10% DMSO

- Centrifuge at  $176 \times g$  for 3 minutes to collect cells.
- Resuspend the cells in pre-cooled freezing medium and adjust the cell density to  $5 \times 10^6$  cells/mL.
- Aliquot 1 mL into each vial.
- Place the vial in a controlled-rate freezing container and store at  $-80^{\circ}\text{C}$  for at least 1 day, then transfer to liquid nitrogen as soon as possible.

## Cell passage

Growth medium: DMEM+10% FBS+1% P.S+15  $\mu\text{g}/\text{mL}$  Blasticidin+0.25  $\mu\text{g}/\text{mL}$  Puromycin

For the first 1 to 2 passages post-resuscitation, use the recovery medium. Once the cells have stabilized, switch to a growth medium.

- Remove and discard culture medium.
- Briefly rinse the cell layer with PBS to remove all traces of serum that contains trypsin inhibitor.
- Add 1.0 mL of 0.25% (w/v) Trypsin-EDTA solution to dish and observe cells under an inverted microscope until cell layer is dispersed (usually within 30 to 60 seconds at  $37^{\circ}\text{C}$ ).
- Note: To avoid clumping do not agitate the cells by hitting or shaking the flask while waiting for the cells to detach. Cells that are difficult to detach may be placed at  $37^{\circ}\text{C}$  to facilitate dispersal.
- Add 2.0 mL of growth medium to mix well and aspirate cells by gently pipetting.
- After centrifugation, resuspend the pellet and add appropriate aliquots of the cell suspension to new culture vessels.
- Incubate cultures at  $37^{\circ}\text{C}$ .

**Subcultivation Ratio: A subcultivation ratio of 1:3 - 1:4 is recommended**

---

**Medium Renewal: Every 2 to 3 days**

## Notes

- a) It is normal to observe a higher number of dead cells immediately after thawing. The condition will improve significantly after adjustment. Once the cells stabilize, the number of dead cells will decrease after subculturing, and the cell growth rate will become stable.

## Sequence

CDH17 Q12864( $\Delta$ ICD)

MILQAHLSLCLLMLYLATGYGQEGKFSGPLKPMTFISIYEGQEPSQIIFQFKANPPAVTFELTGETDNIFVIERE  
 GLLYYNRALDRETRSTHNLQVAALDANGIIVEGVPVITIKVKDINDNRPTFLQSKYEGSVRQNSRPGKPFLYV  
 NATDLDDPATPNGQLYYQIVIQLPMINNVMYFQINNKGTGAISLTREGSQELNPAKNPSYNLVISVKDMGGQSE  
 NSFSDTTSVDIIVTENIWKAPKPVEMVENSTDPHPKITQVRWNDPGAQYSLVDKEKLPRFPFSIDQEGDIYVT  
 QPLDREEKDAYVFYAVAKDEYGVKPLSYPLEIHVKVKDINDNPPTCPSPTVFEVQENERLGNISIGTLTAHDRD  
 EENTANSFLNYRIVEQTPKLPMDGLFLIQTAYAGMLQLAKQSLKKQDTPQYNLTIEVSDKDFKTLCFVQINVIDI  
 NDQIPIFEKSDYGNLTLAEDTNI GSTILTIQATDADEPFTGSSKILYHIIKGDSEGR LGVDTDPHTNTGYVIIKPP  
 LDFETAAVSNI VFKAE NPEPLVFGVKYNASSFAKFTLIVTDVNEAPQFSQHVFQAKVSEDVAIGTKVGNVTAK  
 DPEGLDISYSLRGDTRGWLKIDHVTGEIFSVAPLDREAGSPYRVQVVATEVGGSSLSSVSEFHLILMDVNDNP  
 PRLAKDYTG LFFCHPLSAPGSLIFEATDDDQHLFRGPHFTFSLGSGSLQNDWEVSKINGTHARLSTRHTEFEER  
 EYVVLIRINDGGRPPLEGIVSLPVTFCSCVEGSCFRPAGHQGTGPTVGMVAVGILLTTLLVIGIILAVVFI\*

## Related Products

CDH3	
<a href="#">Cynomolgus_CDH3 CHO-K1 Cell Line</a>	<a href="#">H_CDH3 CHO-K1 Cell Line</a>
<a href="#">H_CDH3 HEK-293 Cell Line</a>	<a href="#">Anti-H_CDH3 hIgG1 Antibody</a>
CDH6	
<a href="#">Cynomolgus_CDH6 CHO-K1 Cell Line</a>	<a href="#">H_CDH6 CHO-K1 Cell Line</a>
<a href="#">H_CDH6 HEK-293 Cell Line</a>	<a href="#">Anti-H_CDH6 hIgG1 Antibody(H01L02)</a>
<a href="#">Anti-CDH6 hIgG1 Reference Antibody (Ralubio)</a>	
CDH17	
<a href="#">Cynomolgus_CDH17 HEK-293 Cell Line</a>	<a href="#">Cynomolgus_CDH17(XP_005563762.1) HEK-293 Cell Line</a>
<a href="#">H_CDH17 CHO-K1 Cell Line</a>	<a href="#">H_CDH17 CT26 Cell Line</a>
<a href="#">H_CDH17 HCT116 Cell Line</a>	<a href="#">H_CDH17 HEK-293 Cell Line</a>
<a href="#">H_CDH17 LLC1 Cell Line</a>	<a href="#">H_CDH17 MC38 Cell Line</a>
<a href="#">H_CDH17 SW480 Cell Line</a>	<a href="#">H_CDH17(<math>\Delta</math>EC1,Flag-EC2-7) HEK-293 Cell Line</a>
<a href="#">H_CDH17(<math>\Delta</math>EC1-2,Flag-EC3-7) HEK-293 Cell Line</a>	<a href="#">H_CDH17(<math>\Delta</math>EC1-3,Flag-EC4-7) HEK-293 Cell Line</a>
<a href="#">H_CDH17(<math>\Delta</math>EC1-4,Flag-EC5-7) HEK-293 Cell Line</a>	<a href="#">H_CDH17(<math>\Delta</math>EC1-5,Flag-EC6-7) HEK-293 Cell Line</a>
<a href="#">H_CDH17(<math>\Delta</math>EC1-6,Flag-EC7) HEK-293 Cell Line</a>	<a href="#">Mouse_CDH17 HEK-293 Cell Line</a>
<a href="#">Rat_CDH17 HEK-293 Cell Line</a>	<a href="#">Rhesus_CDH17 HEK-293 Cell Line</a>

Anti-CDH17 hIgG1 Antibody(BI-905711)	Anti-CDH17 hIgG1 Antibody(VHH1-28BB)
Anti-CDH17 hIgG1 Reference Antibody(BI-905711)	Human CDH17 Protein; His Tag
Mouse CDH17 Protein; His Tag	Biotinylated Human CDH17 Protein; His-Avi Tag
Cynomolgus CDH17 Protein; His Tag	

## Limited Use License Agreement

Genomeditech (Shanghai) Co., Ltd grants to the Licensee all intellectual property rights, exclusive, non-transferable, and non-sublicensable rights of the Licensed Materials; Genomeditech (Shanghai) Co., Ltd will retain ownership of the Licensed Materials, cell line history packages, progeny, and the Licensed Materials including modified materials.

Between Genomeditech (Shanghai) Co., Ltd, and Licensee, Licensee is not permitted to modify cell lines in any way. The Licensee shall not share, distribute, sell, sublicense, or otherwise provide the Licensed Materials, or progenitors to third parties such as laboratories, departments, research institutions, hospitals, universities, or biotechnology companies for use other than for the purpose of outsourcing the Licensee's research.

Please refer to the Genomeditech Cell Line License Agreement for details.

Genomeditech