

# Product Sheet

## H\_αvβ6 LLC1 Cell Line

Catalog number: GM-C32910

Version 3.3.1.241125

<b>Description</b>	H_αvβ6 LLC1 Cell Line is a clonal stable LLC1 cell line that constitutively expresses the human ITGAV and human ITGB6 genes, constructed using lentiviral technolog.
<b>Quantity</b>	5E6 Cells per vial,1 mL
<b>Product Format</b>	1 vial of frozen cells
<b>Shipping</b>	Shipped on dry ice
<b>Storage Conditions</b>	Liquid nitrogen immediately upon receipt
<b>Target</b>	Human_ITGAV & Human_ITGB6
<b>Gene ID/Uniprot ID</b>	P06756-1 & P18564-1
<b>Host Cell</b>	LLC1
<b>Recovery Medium</b>	DMEM+10% FBS+1% P.S
<b>Growth medium</b>	DMEM+10% FBS+1% P.S+200 μg/mL Hygromycin+200 μg/mL Bleomycin
<b>Note</b>	None
<b>Freezing Medium</b>	90% FBS+10% DMSO
<b>Growth properties</b>	Mixed: adherent and suspension
<b>Growth Conditions</b>	37°C, 5% CO <sub>2</sub>
<b>Mycoplasma Testing</b>	The cell line has been screened to confirm the absence of Mycoplasma species.
<b>Safety considerations</b>	Biosafety Level 2
<b>Note</b>	It is recommended to expand the cell culture and store a minimum of 10 vials at an early passage for potential future use.

## Materials

Reagent	Manufacturer/Catalogue No.
DMEM	VivaCell/C3110-0500
Fetal Bovine Serum	Cegrogen biotech/A0500-3010
Pen/Strep	Thermo/15140-122
Bleomycin	Genomeditech/GM-040407
Hygromycin	Genomeditech/GM-040403
Anti- $\alpha$ v hIgG2 Antibody(Abituzumab)	Genomeditech/GM-49298AB
Anti- $\alpha$ v $\beta$ 6 hIgG1 Antibody(m15H3)	Genomeditech/GM-53193AB

## Figures




Figure 1 | H\_ $\alpha$ v $\beta$ 6 LLC1 Cell Line (Cat. GM-C32910) was determined by flow cytometry using Anti- $\alpha$ v hIgG2 Antibody(Abituzumab) (Cat. [GM-49298AB](#)) and Anti- $\alpha$ v $\beta$ 6 hIgG1 Antibody(m15H3) (Cat. [GM-53193AB](#)).

## Cell Recovery

Recovery Medium: DMEM+10% FBS+1% P.S

To insure the highest level of viability, thaw the vial and initiate the culture as soon as possible upon receipt. If upon arrival, continued storage of the frozen culture is necessary, it should be stored in liquid nitrogen vapor phase and not at  $-70^{\circ}\text{C}$ . Storage at  $-70^{\circ}\text{C}$  will result in loss of viability.

- Thaw the vial by gentle agitation in a  $37^{\circ}\text{C}$  water bath. To reduce the possibility of contamination, keep the O-ring and cap out of the water. Thawing should be rapid (approximately 2 - 3 minutes).
- Remove the vial from the water bath as soon as the contents are thawed, and decontaminate by dipping in or spraying with 70% ethanol. All of the operations from this point on should be carried out under strict aseptic conditions.

- c) Transfer the vial contents to a centrifuge tube containing 5.0 mL complete culture medium and spin at approximately 176 x g for 5 minutes. Discard supernatant.
- d) Resuspend cell pellet with the recommended recovery medium. And dispense into appropriate culture dishes.
- e) Incubate the culture at 37°C in a suitable incubator. A 5% CO<sub>2</sub> in air atmosphere is recommended if using the medium described on this product sheet.

## Cell Freezing

Freezing Medium: 90% FBS+10% DMSO

- a) Centrifuge at 176 x g for 3 minutes to collect cells.
- b) Resuspend the cells in pre-cooled freezing medium and adjust the cell density to 5E6 cells/mL.
- c) Aliquot 1 mL into each vial.
- d) Place the vial in a controlled-rate freezing container and store at -80°C for at least 1 day, then transfer to liquid nitrogen as soon as possible.

## Cell passage

Growth medium: DMEM+10% FBS+1% P.S+200 µg/mL Hygromycin+200 µg/mL Bleomycin

For the first 1 to 2 passages post-resuscitation, use the recovery medium. Once the cells have stabilized, switch to a growth medium.

- a) Under normal conditions, these cells exist as both adherent and round suspension cells.
- b) When changing the medium, take care to retain the suspension cells. During passaging, collect both the adherent and suspension cells together before subculturing.
- c) Add 1.0 mL of 0.25% (w/v) Trypsin-EDTA solution to dish and observe cells under an inverted microscope until cell layer is dispersed (usually within 1 to 2 minutes at 37°C).
- d) Note: To avoid clumping do not agitate the cells by hitting or shaking the flask while waiting for the cells to detach. Cells that are difficult to detach may be placed at 37°C to facilitate dispersal.
- e) Add 2.0 mL of growth medium to mix well and aspirate cells by gently pipetting.
- f) After centrifugation, resuspend the pellet and add appropriate aliquots of the cell suspension to new culture vessels.
- g) Incubate cultures at 37°C.

**Subcultivation Ratio: A subcultivation ratio of 1:2 - 1:4 is recommended**

**Medium Renewal: Every 2 to 3 days**

## Notes

- a) After the stabilization of the cell condition, there will be fewer dead cells post-passage, the cell growth rate will tend to stabilize, cell morphology will become uniform, and the cells will appear robust.

## Sequence

### ITGAV [P06756-1](#)

MAFPPRRRLRLGPRGLPLLLSGLLLPLCRAFNLVDSPA EYSGPEGSYFGFAVDFVPSASSRMFLLVGAPKA  
 NTTQPGIVEGGQVLKCDWSSSTRRCQPIEFDATGNRDYAKDDPLEFKSHQWFGASVRSKQDKILACAPLYHW  
 RTEMKQEREPVGTCTFLQDGTKTVEYAPCRSQDIDADGGQFCQGGFSIDFTKADR VLLGGPGSFYWQQLISD  
 QVAEIVSKYDPNVYSIKYNNQLATRTAQAIFFDSSYLGYSVAVGDFNGDGIDDFVSGVPRAARTLGMVYIYDG  
 KNMSSLYNFTGEQMAAYFGFSVAATDINGDDYADVFIGAPLFMDRGS D GKLEVGQVSVSLQRASGDFQTT  
 KLNGFEV FARFGSAIAPLGDLDQDGFNDIAIAAPYGGEDKKGIVYIFNGRSTGLNAVPSQILEGQWAARSMPP  
 SFGYSMKGATDIDKNGYPDLIVGAFGVDRAILYRARPVITVNAGLEVYPSILNQDNKTC SLPGTALKVSCFNV  
 RFCLKADGKGVLPKLNQVVELLDKLGKGAIRRALFLYSRSPSHSKNMTISRGGMLMQCEELIAYLRDESEF  
 RDKLTPITIFMEYRLDYRTAADTTGLQPILNQFTPANISRQA HILLDCGEDNVCKPKLEVSVDSDQKKIYIGDD  
 NPLTLIVKAQNQGE GAYEAELIVSIPLQADFIGVVRNNEALARLSCAFKTENQTRQVVC DLGNPMKAGTQLL  
 AGLRFSVHQQSEMDTSVKFDLQIQSSNLFDKVSPVVSHKVDLAVLA AVEIRGVSSPDHVFLPIPNWEHKENPE  
 TEEDVGPVVQHIYELRNNGPSSFSKAMLHLQWPYKYNNTLLYILHYDIDGPMNCTSDMEINPLRIKISSLQT  
 TEKNDTVAGQGERDHLITKRDLALSEGDIHTL GCGVAQCLKIVCQVGR LDRGKSAILYVKSLWTETFMNKE  
 NQNHSYSLKSSASFNVIEFPYKNLPIEDITNSTLVTNTV TWGIQAPMPVPVWVILAVLAGLLLLAVLVFVMY  
 RMGFFKRV RPPQEEQEREQLQPHENGEGNSET

### ITGB6 [P18564-1](#)

MGIELLCLFFLFLGRNDHVQGGCALGGAETCEDCLLIGPQCAWCAQENFTHPSGVGERCDTPANLLAKGCQL  
 NFIENPVSQVEILKNKPLSVGRQKNSSDIVQIAPQSLILKLRPGGAQTLQVHVRQTEDYPVDLYYLMDSLASM  
 DDDLNTIKELGSRLSKEMSKLTSNFRLLGFGS FVEKPVSPFVKTTPEEIANPCSSIPYFCLPTFGFKHILPLTNDAE  
 RFNEIVKNQKISANIDTPEGGFDAIMQAAVCKEKIGWRNDSLHLLV FVSDADSHFGMDSKLAGIVIPNDGLCH  
 LDSKNEYSMSTVLEYPTIGQLIDKLVQNNVLLIFA VTQEQVHLYENYAKLIPGATVGLLQKDSGNILQLIISAY  
 EELRSEVELEVLGDTEGLNLSFTAICNNGTLFQH QKKCSHMKVGD TASFSVTVNIPH CERRSRHIIKPVGLGD  
 ALELLVSPENCDCQKEVEVNSSKCHHNGSFQCGVCACHPGHMGRCECGEDMLSTDSCKEAPDHPSCSG  
 RGDCYCGQCICHLSPYGNIGPYCQCDNFSCVRHKGLLCGGNGDCDCGECVCRSGWTGEYCNCTTSTDSCV  
 SEDGVLCSGRGDCVCGKCVCTNPGASGPTCERCPTCGDPCNSKRSCIECHLSAAGQAREECVDKCKLAGATI  
 SEEEDFSKDGSVSCSLQGENECLITFLITDNEGKTIHSINEKDCPKPPNIPMIMLGVSLAILLIGVLLCIWKLL  
 VSFHDRKEVAKFEAERSKAKWQTGTNPLYRGSTSTFKNVTYKHREKQKVDLSTDC

## Related Products

TGF-β:GARP:avβ6	
<a href="#">H_GARP Latent TGFB1 Reporter HEK-293 Cell Line</a>	<a href="#">TGF-β Reporter 293 DDX35TM Cell Line</a>
<a href="#">TGF-β Reporter HEK-293 Cell Line</a>	<a href="#">Cynomolgus_αvβ6 HEK-293 Cell Line</a>
<a href="#">H_GARP CHO-K1 Cell Line</a>	<a href="#">H_GARP HEK-293 Cell Line</a>
<a href="#">H_GARP Latent TGF-β1 CHO-K1 Cell Line</a>	<a href="#">H_GARP Latent TGF-β1 HEK-293 Cell Line</a>
<a href="#">H_ITGB6 CHO-K1 Cell Line</a>	<a href="#">H_ITGB6 HEK-293 Cell Line</a>
<a href="#">H_αvβ6 CT26 Cell Line</a>	<a href="#">H_αvβ6 HEK-293 Cell Line</a>
<a href="#">H_αvβ6 MC38 Cell Line</a>	
<a href="#">Anti-GARP-TGF-β1 hIgG4 Antibody(ARGX-115)</a>	<a href="#">Anti-H_ITGB6 hIgG1 Reference Antibody (h2A2)</a>

Anti-ITGB6 hIgG1 Antibody(SGN-B6A)	Anti-TGFB1 hIgG4 Antibody(SRK-181)
Anti- $\alpha$ v hIgG2 Antibody(Abituzumab)	Anti- $\alpha$ v $\beta$ 6 hIgG1 Antibody(m15H3)
Anti-ITGB6-MMAE ADC(Dar4)[SGN-B6A]	
ADC Related Product	
Anti-DXD Mouse IgG1 Antibody (23E21C5)	Anti-DXD Mouse IgG1 Antibody (4A5A12)
Anti-Dxd Mouse IgG2a Antibody (17D6A4)	Anti-Eribulin Mouse IgG2a Antibody (10F8G4)
Anti-MMAE Mouse IgG1 Antibody (11C10E3)	Anti-MMAE Mouse IgG2a Antibody (17A1K11)
Anti-MMAE Mouse IgG2a Antibody (8F6A3)	Mouse anti Human IgG-MMAE(Dar4)
Human IgG1 Isotype-DXD (Dar8)	Human IgG1 Isotype-Eribulin (Dar4)
Human IgG1 Isotype-MMAE (Dar4)	
Recombinant DT3C Protein	

## Limited Use License Agreement

Genomeditech (Shanghai) Co., Ltd grants to the Licensee all intellectual property rights, exclusive, non-transferable, and non-sublicensable rights of the Licensed Materials; Genomeditech (Shanghai) Co., Ltd will retain ownership of the Licensed Materials, cell line history packages, progeny, and the Licensed Materials including modified materials.

Between Genomeditech (Shanghai) Co., Ltd, and Licensee, Licensee is not permitted to modify cell lines in any way. The Licensee shall not share, distribute, sell, sublicense, or otherwise provide the Licensed Materials, or progenitors to third parties such as laboratories, departments, research institutions, hospitals, universities, or biotechnology companies for use other than for the purpose of outsourcing the Licensee's research.

Please refer to the Genomeditech Cell Line License Agreement for details.