

# Product Sheet

## H\_ACVR2B HEK-293(ACVR2A KO) Cell Line

Catalog number: GM-C34942

Version 3.3.1.241204

<b>Description</b>	H_ACVR2B HEK-293(ACVR2A KO) Cell Line is a clonal stable HEK-293 cell line that continuously expresses human ACVR2B. It is constructed using lentiviral technology, based on the knockout of human ACVR2A.
<b>Quantity</b>	5E6 Cells per vial, 1 mL
<b>Product Format</b>	1 vial of frozen cells
<b>Shipping</b>	Shipped on dry ice
<b>Storage Conditions</b>	Liquid nitrogen immediately upon receipt
<b>Target</b>	Human_ACVR2B
<b>Gene ID/Uniprot ID</b>	Q13705-1
<b>Host Cell</b>	HEK-293
<b>Recovery Medium</b>	DMEM+10% FBS+1% P.S
<b>Growth medium</b>	DMEM+10% FBS+1% P.S+125 µg/mL Hygromycin
<b>Note</b>	The cells are very sensitive to antibiotics and should be cultured using the cell growth medium provided by Genomeditech.
<b>Freezing Medium</b>	90% FBS+10% DMSO
<b>Growth properties</b>	Adherent
<b>Growth Conditions</b>	37°C, 5% CO <sub>2</sub>
<b>Mycoplasma Testing</b>	The cell line has been screened to confirm the absence of Mycoplasma species.
<b>Safety considerations</b>	Biosafety Level 2
<b>Note</b>	It is recommended to expand the cell culture and store a minimum of 10 vials at an early passage for potential future use.

## Materials

Reagent	Manufacturer/Catalogue No.
DMEM	Gibco/C11995500BT
Fetal Bovine Serum	Cegrogen biotech/A0500-3010
Pen/Strep	Thermo/15140-122
Hygromycin	Genomeditech/GM-040403
Anti-ACVR2B hIgG1 Antibody(Bimagrumab)	Genomeditech/GM-51148AB
Anti-ACVR2B hIgG1 Antibody(Fab-17G05)	Genomeditech/GM-82350AB

## Figures




Figure 1 | H\_ACVR2B HEK-293(ACVR2A KO) Cell Line (Cat. GM-C34942) was determined by flow cytometry using Anti-ACVR2B hIgG1 Antibody(Fab-17G05) (Cat. [GM-82350AB](#)).




Figure 2 | H\_ACVR2B HEK-293(ACVR2A KO) Cell Line (Cat. GM-C34942) was determined by flow cytometry using Anti-ACVR2B hIgG1 Antibody(Bimagrumab) (Cat. [GM-51148AB](#)).




Figure 3 | H\_ACVR2A HEK-293(ACVR2B KO) Cell Line (Cat. GM-C34941) was determined by flow cytometry using Anti-ACVR2B hIgG1 Antibody(Fab-17G05) (Cat. [GM-82350AB](#)).



Figure 4 | The Sanger sequencing of the parental cells of the H\_ACVR2B HEK-293 (ACVR2A KO) cell line showed successful knockout of ACVR2A.




Figure 5 | The passage stability of the H\_ACVR2B HEK-293(ACVR2A KO) Cell Line (Cat. GM-C34942) was determined by flow cytometry using Anti-ACVR2B hIgG1 Antibody(Fab-17G05) (Cat. [GM-82350AB](#)).

## Cell Recovery

Recovery Medium: DMEM+10% FBS+1% P.S

To insure the highest level of viability, thaw the vial and initiate the culture as soon as possible upon receipt. If upon arrival, continued storage of the frozen culture is necessary, it should be stored in liquid nitrogen vapor phase and not at  $-70^{\circ}\text{C}$ . Storage at  $-70^{\circ}\text{C}$  will result in loss of viability.

- a) Thaw the vial by gentle agitation in a  $37^{\circ}\text{C}$  water bath. To reduce the possibility of contamination, keep the O-ring and cap out of the water. Thawing should be rapid (approximately 2 - 3 minutes).
- b) Remove the vial from the water bath as soon as the contents are thawed, and decontaminate by dipping in or spraying with 70% ethanol. All of the operations from this point on should be carried out under strict aseptic conditions.
- c) Transfer the vial contents to a centrifuge tube containing 5.0 mL complete culture medium and spin at approximately  $176 \times g$  for 5 minutes. Discard supernatant.
- d) Resuspend cell pellet with the recommended recovery medium. And dispense into appropriate culture dishes.
- e) Incubate the culture at  $37^{\circ}\text{C}$  in a suitable incubator. A 5%  $\text{CO}_2$  in air atmosphere is recommended if using the medium described on this product sheet.

## Cell Freezing

Freezing Medium: 90% FBS+10% DMSO

- a) Centrifuge at  $176 \times g$  for 3 minutes to collect cells.
- b) Resuspend the cells in pre-cooled freezing medium and adjust the cell density to  $5 \times 10^6$  cells/mL.
- c) Aliquot 1 mL into each vial.
- d) Place the vial in a controlled-rate freezing container and store at  $-80^{\circ}\text{C}$  for at least 1 day, then transfer to liquid nitrogen as soon as possible.

## Cell passage

Growth medium: DMEM+10% FBS+1% P.S+125  $\mu\text{g}/\text{mL}$  Hygromycin

For the first 1 to 2 passages post-resuscitation, use the recovery medium. Once the cells have stabilized, switch to a growth medium.

- a) Subculturing is necessary when the cell density reaches 80%. It is recommended to perform subculturing at a ratio of 1:3 to 1:4 every 2-3 days. Ensure that the density does not exceed 80%, as overcrowding can lead to reduced viability due to compression.
- b) Remove and discard culture medium.
- c) Briefly rinse the cell layer with PBS to remove all traces of serum that contains trypsin inhibitor.
- d) Add 1.0 mL of 0.25% (w/v) Trypsin-EDTA solution to dish and observe cells under an inverted microscope until cell layer is dispersed (usually within 30 to 60 seconds at  $37^{\circ}\text{C}$ ).
- e) Note: To avoid clumping do not agitate the cells by hitting or shaking the flask while waiting for the cells to detach. Cells that are difficult to detach may be placed at  $37^{\circ}\text{C}$  to facilitate dispersal.
- f) Add 2.0 mL of growth medium to mix well and aspirate cells by gently pipetting.
- g) After centrifugation, resuspend the pellet and add appropriate aliquots of the cell suspension to new culture vessels.
- h) Incubate cultures at  $37^{\circ}\text{C}$ .

**Subcultivation Ratio: A subcultivation ratio of 1:3 - 1:4 is recommended**

**Medium Renewal: Every 2 to 3 days**

## Notes

- Upon initial thawing, a higher number of dead cells is observed, which is a normal phenomenon. Significant improvement is seen after adaptation. Once the cells reach a stable state, the number of dead cells decreases after subculturing and the cell growth rate becomes stable.
- Ensure that the cell density does not exceed 80%, as overcrowding may lead to reduced viability due to compression.

## Sequence

ACVR2B Q13705-1

MTAPWVALALLWGSLCAGSGRGEAETRECIYYNANWELERTNQSGLERCEGEQDKRLHCYASWRNSSGTIE  
 LVKKGCVLDDFNCYDRQECVATEENPQVYFCCCEGNFCNERFTHLPEAGGPEVTYEPPPTAPLLTLVLA  
 LPIGGLSLIVLLAFWVYRHRKPPYGHVDIHEDPGPPPSPLVGLKPLQLEIKARGRFGCVWKAQLMNDFVA  
 VKIFPLQDKQSWQSEREIFSTPGMKHENLLQFIAAEKRGSNLEVELWLITAFHDKGSLTDYLGKNIITWNE  
 LCHVAETMSRGLSYLHEDVPCWCRGEGHKSIAHRDFKSKNVLLKSDLTAVLADFGLA VRFEPGKPPGDTHGQV  
 GTRRYMAPEVLEGAINFQRDAFLRIDMYAMGLVLWELVSRCKAADGPVDEYMLPFEEEEIGQHP  
 SLEELQEVVVHKKMRPTIKDHWLKHPLAQLCVTIEECWDHDAEARLSAGCVEERVSLIRRSVNGTTS  
 DCLVSLVTSVTNVDLPPKESSI

## Related Products

GCGR	
<a href="#">H_GCGR Reporter CHO-K1 Cell Line</a>	<a href="#">H_GCGR Reporter HEK-293 Cell Line</a>
<a href="#">H_GCGR CHO-K1 Cell Line</a>	<a href="#">H_GCGR HEK-293 Cell Line</a>
<a href="#">Mouse_GCGR HEK-293 Cell Line</a>	
<a href="#">Anti-H_GCGR hIgG2 Antibody(volagidemab)</a>	
GLP1R	
<a href="#">H_GLP1R Reporter CHO-K1 Cell Line</a>	<a href="#">H_GLP1R Reporter HEK-293 Cell Line</a>
<a href="#">H_GLP1R Reporter HEK-293 DDX35TM Cell Line</a>	<a href="#">Cynomolgus_GLP1R HEK-293 Cell Line</a>
<a href="#">H_GLP1R CHO-K1 Cell Line</a>	<a href="#">H_GLP1R HEK-293 Cell Line</a>
<a href="#">Mouse_GLP1R HEK-293 Cell Line</a>	
<a href="#">Anti-GLP1R hIgG1 Antibody(mAb-36986)</a>	<a href="#">Anti-H_GLP1R hIgG1 Antibody(glutazumab)</a>
FGF21:FGFR	
<a href="#">H_FGF21 Reporter HEK-293 Cell Line</a>	
CALCA(CGRP): CALCRL RAMP	
<a href="#">H_CALCRL RAMP1 Reporter HEK-293 Cell Line</a>	<a href="#">Cynomolgus_CALCRL RAMP1 HEK-293 Cell Line</a>
<a href="#">H_CALCRL RAMP1 CHO-K1 Cell Line</a>	<a href="#">H_CALCRL RAMP1 HEK-293 Cell Line</a>
<a href="#">Anti-CALCRL RAMP1 hIgG2 Antibody(Erenumab)</a>	
GIP:GIPR	

H_GIPR Reporter CHO-K1 Cell Line	H_GIPR Reporter HEK-293 Cell Line
H_GIPR Reporter HEK-293 DDX35TM Cell Line	Cynomolgus_GIPR HEK-293 Cell Line
H_GIPR CHO-K1 Cell Line	H_GIPR HEK-293 Cell Line
Mouse_GIPR HEK-293 Cell Line	
Anti-H_GIPR hIgG1 Antibody(AMG-133)	
<b>ACVR2A: ACTRIIB: Active A</b>	
ACVR2A KO HEK-293 Cell Line	Activin A Reporter Cell Line
H_ACVR2A Reporter Cell Line	H_ACVR2B Reporter Cell Line
ACVR2B KO HEK-293 Cell Line	H_ACVR2A HEK-293(ACVR2B KO) Cell Line
Anti-ACVR2B hIgG1 Antibody(Bimagrumab)	Anti-ACVR2B hIgG1 Antibody(Fab-17G05)
Anti-ACVR2B mIgG2a Antibody(Bimagrumab)	Anti-H_ACVR2B hIgG1 Reference Antibody(Bimbio)
Biotinylated Human ACVR2A Protein; His-Avi Tag	Biotinylated Human ACVR2B Protein; His-Avi Tag
Biotinylated Mouse ACVR2A Protein; His-Avi Tag	Biotinylated Mouse ACVR2B Protein; His-Avi Tag
Human Activin A Protein; His Tag	Human Activin B Protein; His Tag
Human ACVR2A Protein; hFc Tag	Human ACVR2A Protein; His Tag
Human ACVR2B Protein; hFc Tag	Human ACVR2B Protein; His Tag
Mouse ACVR2B Protein; His Tag	
<b>AMY: CALCR RAMP</b>	
H_CALCR RAMP3(AMY3) Reporter CHO-K1 Cell Line	H_CALCR Reporter CHO-K1 Cell Line

## Limited Use License Agreement

Genomeditech (Shanghai) Co., Ltd grants to the Licensee all intellectual property rights, exclusive, non-transferable, and non-sublicensable rights of the Licensed Materials; Genomeditech (Shanghai) Co., Ltd will retain ownership of the Licensed Materials, cell line history packages, progeny, and the Licensed Materials including modified materials.

Between Genomeditech (Shanghai) Co., Ltd, and Licensee, Licensee is not permitted to modify cell lines in any way. The Licensee shall not share, distribute, sell, sublicense, or otherwise provide the Licensed Materials, or progenitors to third parties such as laboratories, departments, research institutions, hospitals, universities, or biotechnology companies for use other than for the purpose of outsourcing the Licensee's research.

Please refer to the Genomeditech Cell Line License Agreement for details.