

# Product Sheet

## H\_DLL3 B16-F10 Cell Line

Catalog number: GM-C35055

Version 3.3.1.250619

<b>Description</b>	H_DLL3 B16-F10 Cell Line is a clonal stable B16-F10 cell line that constitutively expresses the human DLL3 gene, constructed using lentiviral technology.
<b>Quantity</b>	5E6 Cells per vial, 1 mL
<b>Product Format</b>	3 vials of frozen cells
<b>Shipping</b>	Shipped on dry ice
<b>Storage Conditions</b>	Liquid nitrogen immediately upon receipt
<b>Target</b>	Human_DLL3
<b>Gene ID/Uniprot ID</b>	Q9NYJ7(AA Ala 27 - Leu 492)
<b>Host Cell</b>	B16-F10
<b>Recovery Medium</b>	DMEM+10% FBS+1% P.S
<b>Growth medium</b>	DMEM+10% FBS+1% P.S+0.5 µg/mL Puromycin
<b>Note</b>	None
<b>Freezing Medium</b>	90% FBS+10% DMSO
<b>Growth properties</b>	Adherent
<b>Growth Conditions</b>	37°C, 5% CO <sub>2</sub>
<b>Mycoplasma Testing</b>	The cell line has been screened to confirm the absence of Mycoplasma species.
<b>Safety considerations</b>	Biosafety Level 2
<b>Note</b>	It is recommended to expand the cell culture and store a minimum of 10 vials at an early passage for potential future use.

## Materials

Reagent	Manufacturer/Catalogue No.
DMEM	Gibco/C11995500BT
Fetal Bovine Serum	Cegrogen biotech/A0500-3010
Pen/Strep	Thermo/15140-122
Puromycin	Genomeditech/ <a href="#">GM-040401</a>
Anti-H_DLL3 hIgG1 Antibody(Rovalpituzumab)	Genomeditech/ <a href="#">GM-26560AB</a>

## Figures




Figure 1 | H\_DLL3 B16-F10 Cell Line (Cat. GM-C35055) was determined by flow cytometry using Anti-H\_DLL3 hIgG1 Antibody(Rovalpituzumab) (Cat. [GM-26560AB](#)).

## Cell Recovery

Recovery Medium: DMEM+10% FBS+1% P.S

To insure the highest level of viability, thaw the vial and initiate the culture as soon as possible upon receipt. If upon arrival, continued storage of the frozen culture is necessary, it should be stored in liquid nitrogen vapor phase and not at -70°C. Storage at -70°C will result in loss of viability.

- Thaw the vial by gentle agitation in a 37°C water bath. To reduce the possibility of contamination, keep the O-ring and cap out of the water. Thawing should be rapid (approximately 2 - 3 minutes).
- Remove the vial from the water bath as soon as the contents are thawed, and decontaminate by dipping in or spraying with 70% ethanol. All of the operations from this point on should be carried out under strict aseptic conditions.
- Transfer the vial contents to a centrifuge tube containing 5.0 mL complete culture medium and spin at approximately 176 x g for 5 minutes. Discard supernatant.
- Resuspend cell pellet with the recommended recovery medium. And dispense into appropriate culture dishes.

- e) Incubate the culture at 37°C in a suitable incubator. A 5% CO<sub>2</sub> in air atmosphere is recommended if using the medium described on this product sheet.

## Cell Freezing

Freezing Medium: 90% FBS+10% DMSO

- Centrifuge at 176 x g for 3 minutes to collect cells.
- Resuspend the cells in pre-cooled freezing medium and adjust the cell density to 5E6 cells/mL.
- Aliquot 1 mL into each vial.
- Place the vial in a controlled-rate freezing container and store at -80°C for at least 1 day, then transfer to liquid nitrogen as soon as possible.

## Cell passage

Growth medium: DMEM+10% FBS+1% P.S+0.5 µg/mL Puromycin

For the first 1 to 2 passages post-resuscitation, use the recovery medium. Once the cells have stabilized, switch to a growth medium.

- Remove and discard culture medium.
- Briefly rinse the cell layer with PBS to remove all traces of serum that contains trypsin inhibitor.
- Add 1.0 mL of 0.25% (w/v) Trypsin-EDTA solution to dish and observe cells under an inverted microscope until cell layer is dispersed (usually within 30 to 60 seconds at 37°C).
- Note: To avoid clumping do not agitate the cells by hitting or shaking the flask while waiting for the cells to detach. Cells that are difficult to detach may be placed at 37°C to facilitate dispersal.
- Add 2.0 mL of growth medium to mix well and aspirate cells by gently pipetting.
- After centrifugation, resuspend the pellet and add appropriate aliquots of the cell suspension to new culture vessels.
- Incubate cultures at 37°C.

**Subcultivation Ratio: A subcultivation ratio of 1:3 - 1:4 is recommended**

**Medium Renewal: Every 2 to 3 days**

## Notes

- If small black dots appear inside the cells, particles are present in the intercellular spaces, or a few dead cells are floating in the medium during the cultivation process, these are all common phenomena in cell culture and do not affect cell proliferation. The cell pellet after centrifugation may also appear black.

## Sequence

DLL3 Q9NYJ7(ECD)

AGVFELQIHSFGPGPGAPRSPCSARLPCRLFFRVCLKPGLSEEAESPALGAALSARGPVYTEQPGAPAPD  
LPLPDGLLQVPFRDAWPGTFSFIETWREELGDQIGGPAWSLLARVAGRRRLAAGGPWARDIQRAGAWELRF  
SYRARCEPPAVGTACTRLCRPRSAPSRCGPGLRPCAPLEDECEAPLVCRAGCSPEHGFCEQPGECRCLEGWTG

PLCTVPVSTSSCLSPRGPSSATTGCLVPGPGPCDGNPCANGGSCSETPRSFECTCPRGFYGLRCEVSGVTCADG  
PCFNGLCVGGADPDSAYICHCPPGFQGSNCEKRVDRCSLQPCRNGGLCLDLGHALRCRCRAGFAGPRCEH  
DLDDCAGRACANGGTCVEGGGAHRCSCALGFGGRDCRERADPCAARPCAHHGRCYAHFSGLVACAPGY  
MGARCEFPVHPDGASALPAAPPGLRPGDPQRYL

## Related Products

DLL1	
<a href="#">H_DLL1 CHO-K1 Cell Line</a>	
<a href="#">Anti-DLL1 hIgG1 Antibody(pidilizumab)</a>	
CD3	
<a href="#">H_CD3D CD3E KO Jurkat Cell Line</a>	<a href="#">Jurkat CD3-BsAb Reporter Cell Line</a>
<a href="#">Cynomolgus_CD3 HEK-293 Cell Line</a>	<a href="#">Cynomolgus_CD3E(Membrane Bound ECD) CHO-K1 Cell Line</a>
<a href="#">H_CD3 CHO-K1 Cell Line</a>	<a href="#">H_CD3 HEK-293 Cell Line</a>
<a href="#">H_CD3E(Membrane Bound ECD) CHO-K1 Cell Line</a>	<a href="#">Mouse_CD3 HEK-293 Cell Line</a>
<a href="#">Anti-CD3 epsilon hIgG1 Antibody [OKT-3 (muromonab)]</a>	<a href="#">Anti-CD3 hIgG1 Antibody(CH2527)</a>
<a href="#">Anti-mouse CD3ε mIgG2a Antibody(145-2C11)</a>	
DLL4	
<a href="#">Cynomolgus_DLL4 CHO-K1 Cell Line</a>	<a href="#">H_DLL4 CHO-K1 Cell Line</a>
<a href="#">H_DLL4 HEK-293 Cell Line</a>	
<a href="#">Anti-DLL4 hIgG1 Antibody(MLCK-2(ABL-001))</a>	<a href="#">Anti-DLL4 hIgG2 Antibody(Navicixizumab)</a>
DLL3	
<a href="#">Cynomolgus_DLL3 CHO-K1 Cell Line</a>	<a href="#">H_DLL3 CHO-K1 Cell Line</a>
<a href="#">H_DLL3 CT26 Cell Line</a>	<a href="#">H_DLL3 HEK-293 Cell Line</a>
<a href="#">Anti-DLL3 hIgG1 Reference Antibody(Rovabio)</a>	<a href="#">Anti-H_DLL3 hIgG1 Antibody(Rovalpituzumab)</a>
ADC Related Product	
<a href="#">Anti-DXD Mouse IgG1 Antibody (23E21C5)</a>	<a href="#">Anti-DXD Mouse IgG1 Antibody (4A5A12)</a>
<a href="#">Anti-Dxd Mouse IgG2a Antibody (17D6A4)</a>	<a href="#">Anti-Eribulin Mouse IgG2a Antibody (10F8G4)</a>
<a href="#">Anti-MMAE Mouse IgG1 Antibody (11C10E3)</a>	<a href="#">Anti-MMAE Mouse IgG2a Antibody (17A1K11)</a>
<a href="#">Anti-MMAE Mouse IgG2a Antibody (8F6A3)</a>	<a href="#">Mouse anti Human IgG1-MMAE(Dar4)</a>
<a href="#">Human IgG1 Isotype-DXD (Dar8)</a>	<a href="#">Human IgG1 Isotype-Eribulin (Dar4)</a>
<a href="#">Human IgG1 Isotype-MMAE (Dar4)</a>	
<a href="#">Recombinant DT3C Protein</a>	

## Limited Use License Agreement

Genomeditech (Shanghai) Co., Ltd grants to the Licensee all intellectual property rights, exclusive, non-transferable, and non-sublicensable rights of the Licensed Materials; Genomeditech (Shanghai) Co., Ltd will retain ownership of the Licensed Materials, cell line history packages, progeny, and the Licensed Materials including modified materials.

Between Genomeditech (Shanghai) Co., Ltd, and Licensee, Licensee is not permitted to modify cell lines in any way. The Licensee shall not share, distribute, sell, sublicense, or otherwise provide the Licensed Materials, or progenitors to third parties such as laboratories, departments, research institutions, hospitals, universities, or biotechnology companies for use other than for the purpose of outsourcing the Licensee's research.

Please refer to the Genomeditech Cell Line License Agreement for details.

Genomeditech