

Product Sheet

H_SLC39A6 CHO-K1 Cell Line

Catalog number: GM-C13341

Version 3.3.1.250815

Description	H_SLC39A6 CHO-K1 Cell Line is a clonal stable CHO-K1 cell line constitutively expressing human SLC39A6.
Quantity	5E6 Cells per vial, 1 mL
Product Format	1 vial of frozen cells
Shipping	Shipped on dry ice
Storage Conditions	Liquid nitrogen immediately upon receipt
Target	Human_SLC39A6
Gene ID/Uniprot ID	Q13433-1
Host Cell	CHO-K1
Recovery Medium	F12K+10% FBS+1% P.S
Growth medium	F12K+10% FBS+1% P.S+4 µg/mL Puromycin
Note	None
Freezing Medium	90% FBS+10% DMSO
Growth properties	Adherent
Growth Conditions	37°C, 5% CO ₂
Mycoplasma Testing	The cell line has been screened to confirm the absence of Mycoplasma species.
Safety considerations	Biosafety Level 2
Note	It is recommended to expand the cell culture and store a minimum of 10 vials at an early passage for potential future use.

Materials

Reagent	Manufacturer/Catalogue No.
F12K	BOSTER/PYG0036
Fetal Bovine Serum	ExCell/FSP500
Puromycin	Genomeditech/ GM-040401
Anti-H_SLC39A6 hIgG1 Antibody(Ladiratuzumab)	Genomeditech/ GM-26839AB

Figures




Figure 1 | H_SLC39A6 CHO-K1 Cell Line was determined by flow cytometry using Anti-H_SLC39A6 hIgG1 Antibody(Ladiratuzumab) (Genomeditech/[GM-26839AB](#)).

Cell Recovery

Recovery Medium: F12K+10% FBS+1% P.S

To insure the highest level of viability, thaw the vial and initiate the culture as soon as possible upon receipt. If upon arrival, continued storage of the frozen culture is necessary, it should be stored in liquid nitrogen vapor phase and not at -70°C. Storage at -70°C will result in loss of viability.

- Thaw the vial by gentle agitation in a 37°C water bath. To reduce the possibility of contamination, keep the O-ring and cap out of the water. Thawing should be rapid (approximately 2 - 3 minutes).
- Remove the vial from the water bath as soon as the contents are thawed, and decontaminate by dipping in or spraying with 70% ethanol. All of the operations from this point on should be carried out under strict aseptic conditions.
- Transfer the vial contents to a centrifuge tube containing 5.0 mL complete culture medium and spin at approximately 176 x g for 5 minutes. Discard supernatant.
- Resuspend cell pellet with the recommended recovery medium. And dispense into appropriate culture dishes.

- e) Incubate the culture at 37°C in a suitable incubator. A 5% CO₂ in air atmosphere is recommended if using the medium described on this product sheet.

Cell Freezing

Freezing Medium: 90% FBS+10% DMSO

- Centrifuge at 176 x g for 3 minutes to collect cells.
- Resuspend the cells in pre-cooled freezing medium and adjust the cell density to 5E6 cells/mL.
- Aliquot 1 mL into each vial.
- Place the vial in a controlled-rate freezing container and store at -80°C for at least 1 day, then transfer to liquid nitrogen as soon as possible.

Cell passage

Growth medium: F12K+10% FBS+1% P.S+4 µg/mL Puromycin

For the first 1 to 2 passages post-resuscitation, use the recovery medium. Once the cells have stabilized, switch to a growth medium.

- Remove and discard culture medium.
- Briefly rinse the cell layer with PBS to remove all traces of serum that contains trypsin inhibitor.
- Add 1.0 mL of 0.25% (w/v) Trypsin-EDTA solution to dish and observe cells under an inverted microscope until cell layer is dispersed (usually within 2 to 3 minutes at 37°C).
- Note: To avoid clumping do not agitate the cells by hitting or shaking the flask while waiting for the cells to detach. Cells that are difficult to detach may be placed at 37°C to facilitate dispersal.
- Add 2.0 mL of growth medium to mix well and aspirate cells by gently pipetting.
- After centrifugation, resuspend the pellet and add appropriate aliquots of the cell suspension to new culture vessels.
- Incubate cultures at 37°C.

Subcultivation Ratio: A subcultivation ratio of 1:4 - 1:5 is recommended

Medium Renewal: Every 2 to 3 days

Notes

- After the stabilization of the cell condition, there will be fewer dead cells post-passage, the cell growth rate will tend to stabilize, cell morphology will become uniform, and the cells will appear robust.

Sequence

SLC39A6 Q13433-1

MARKLSVILITFALSVTNPLHELKAAAFQTTEKISPWESGINVDLAISTRQYHLQQLFYRYGENNSLSVEG
FRKLLQNIQIDKIKRIHHHDHSDHEHHSDHERHSDHEHHSEHEHSDHDHSHHNHAASGKNKRKALC
PDHSDSDSSGKDPRNSQKGGAHRPEHASGRNRVKSVSASEVTSTVYNTVSEGTHTFLETIETPRPGKLFPKDVS
SSTPPSVTSKSRVSRLAGRKTNESVSEPRKGFMYSRNTNENPQECFNASKLLTSHGMGIQVPLNATEFNLYLCP

AIINQIDARSCLHTSEKKAIEIPPKTYSLQIAWVGGFIAISIISFLSLLGVILVPLMNRVFFKFLLSFLVALAVGTL
 GDAFLHLLPHSHASHHHSHSHEEPAMEMKRGPLFSHLSSQNIEESAYFDSTWKGLTALGGLYFMFLVEHVL
 LIKQFKDKKKKNQKKPENDDDVEIKKQLSKYESQLSTNEEKVDTDDRTGYLRADSQEPHFDSQQPAVLEE
 EEVMIAHAHPQEVYNEYVPRGCKNKCHSHFHDTLGQSDDLIHHDYHHILHHHHHQNHHPHSHSQRYRSR
 ELKDAGVATLAWMVIMGDGLHNFSDGLAIGAAFTEGLSSGLSTSVAVFCHELPHLGDFAVLLKAGMTVKQ
 AVLYNALSAMLAYLGMATGIFIGHYAENVSMWIFALTAGLFMYVALVDMVPEMLHNDASDHGCSRWGYF
 FLQNAGMLLGFGIMLLISIFEHKIVFRINF

Related Products

HER3(ERBB3)	
Cynomolgus_ERBB3(HER3) CHO-K1 Cell Line	Cynomolgus_ERBB3(HER3) HEK-293 Cell Line
H_ERBB3(HER3) CHO-K1 Cell Line	H_ERBB3(HER3) HEK-293 Cell Line
H_ERBB3(HER3) MC38 Cell Line	Mouse_HER3(ERBB3) CHO-K1 Cell Line
Anti-ERBB3(HER3) hIgG1 Reference Antibody(Patibio)	Anti-H_ERBB3(HER3) hIgG1 Antibody(Barecetamab)
Biotinylated Human HER3 Protein; His-Avi Tag	Human HER3 Protein; His Tag
Mouse HER3 Protein; His Tag	
NECTIN4	
H_NECTIN4(nectin-4) CHO-K1 Cell Line	Cynomolgus_Nectin4 CHO-K1 Cell Line
H_NECTIN4 CT26 Cell Line	H_NECTIN4 HEK-293 Cell Line
H_NECTIN4 LLC1 Cell Line	H_NECTIN4 MC38 Cell Line
Anti-H_Nectin4 hIgG1 Antibody(Enfortumab)	Anti-Nectin 4-MMAE(Dar4)[Enfortumab vedotin]
Anti-Nectin4 hIgG1 Reference Antibody (Enfobio)	
Biotinylated Cynomolgus Nectin-4 Protein; His-Avi Tag	Biotinylated Human Nectin-4 Protein; His-Avi Tag
Biotinylated Mouse Nectin-4 Protein; His-Avi Tag	Cynomolgus Nectin-4 Protein; His Tag
Human Nectin-4 Protein; His Tag	
SLC39A6 (LIV1)	
Cynomolgus_SLC39A6 CHO-K1 Cell Line	H_SLC39A6 HEK-293 Cell Line
H_SLC39A6 LLC1 Cell Line	H_SLC39A6 MC38 Cell Line
Anti-H_SLC39A6 hIgG1 Antibody(Ladiratuzumab)	Anti-SLC39A6 hIgG1 Reference Antibody (Ladbio)
Anti-SLC39A6-MMAE ADC(Dar4)[Ladiratuzumab vedotin]	
HER2(ERBB2)	
H_HER2 HER4 Reporter HEK-293 Cell Line	Cynomolgus_HER2(ERBB2) CHO-K1 Cell Line
H_HER2 EMT6 Cell Line	H_HER2 HER3 MC38 Cell Line
H_HER2 MCF-7 Cell Line	H_HER2(ERBB2) CHO-K1 Cell Line
H_HER2(ERBB2) CT26 Cell Line	H_HER2(ERBB2) LLC1 Cell Line
H_HER2(ERBB2) MC38 Cell Line	
Anti-H_HER2 hIgG1 Antibody(Margetuximab)	Anti-HER2 hIgG1 Reference Antibody(Marbio)
Anti-HER2 hIgG1 Reference Antibody(Trasbio)	Anti-HER2-DM1 ADC(Dar4)[Trastuzumab emtansine,T-DM1]
Anti-HER2-DXD ADC(Dar8)[Trastuzumab Deruxtecan]	
Cynomolgus HER2 Protein; His Tag	Human HER2 Protein; His Tag

ADC Related Product	
Anti-DXD Mouse IgG1 Antibody (23E21C5)	Anti-DXD Mouse IgG1 Antibody (4A5A12)
Anti-Dxd Mouse IgG2a Antibody (17D6A4)	Anti-Eribulin Mouse IgG2a Antibody (10F8G4)
Anti-MMAE Mouse IgG1 Antibody (11C10E3)	Anti-MMAE Mouse IgG2a Antibody (17A1K11)
Anti-MMAE Mouse IgG2a Antibody (8F6A3)	Mouse anti Human IgG1-MMAE(Dar4)
Human IgG1 Isotype-DXD (Dar8)	Human IgG1 Isotype-Eribulin (Dar4)
Human IgG1 Isotype-MMAE (Dar4)	
Recombinant DT3C Protein	

License Agreement:

By purchasing and using this cell line product, the user voluntarily agrees to accept and abide by the following policies:

- This cell line product is restricted to research use only and shall not be used for any commercial purposes.
- This product is strictly prohibited from being used in the diagnosis or treatment of human or animal diseases, and shall not be directly used in experiments involving humans.
- Users and their contractors engaged for their benefit may use this material and its derivatives only within the agreed research scope; modification of the material is not permitted, nor may it be distributed, sold, transferred, or otherwise provided to any other entity (including affiliates).
- If use beyond the above scope is required, prior written permission from Genomeditech (Shanghai) Co.,Ltd. must be obtained. For details, please contact Genomeditech (Shanghai) Co.,Ltd.