

# Product Sheet

## Cynomolgus\_GUCY2C HEK-293 Cell Line

Catalog number: GM-C39498

Version 3.3.1.260603

<b>Description</b>	Cynomolgus_GUCY2C HEK-293 Cell Line is a clonal stable HEK-293 cell line that constitutively expresses the cynomolgus GUCY2C gene, constructed using lentiviral technology.
<b>Quantity</b>	5E6 Cells per vial, 1 mL
<b>Product Format</b>	1 vial of frozen cells
<b>Shipping</b>	Shipped on dry ice
<b>Storage Conditions</b>	Liquid nitrogen immediately upon receipt
<b>Target</b>	Cynomolgus_GUCY2C
<b>Gene ID/Uniprot ID</b>	G7PJX5
<b>Host Cell</b>	HEK-293
<b>Recovery Medium</b>	DMEM+10% FBS+1% P.S
<b>Growth medium</b>	DMEM+10% FBS+1% P.S+0.75 µg/mL Puromycin
<b>Note</b>	None
<b>Freezing Medium</b>	90% FBS+10% DMSO
<b>Growth properties</b>	Adherent
<b>Growth Conditions</b>	37°C, 5% CO <sub>2</sub>
<b>Mycoplasma Testing</b>	The cell line has been screened to confirm the absence of Mycoplasma species.
<b>Safety considerations</b>	Biosafety Level 2
<b>Note</b>	It is recommended to expand the cell culture and store a minimum of 10 vials at an early passage for potential future use.

## Materials

Reagent	Manufacturer/Catalogue No.
DMEM	Gibco/C11995500BT
Fetal Bovine Serum	ExCell/FSP500
Pen/Strep	Thermo/15140-122
Puromycin	Genomeditech/GM-040401
Anti-H_GUCY2C hIgG1 Antibody(Indusatumab)	Genomeditech/GM-28860AB

## Figures



Figure 1 | Cynomolgus\_GUCY2C HEK-293 Cell Line (Cat. GM-C39498) was determined by flow cytometry using Anti-H\_GUCY2C hIgG1 Antibody(Indusatumab) (Cat. [GM-28860AB](#)).

## Cell Recovery

Recovery Medium: DMEM+10% FBS+1% P.S

To insure the highest level of viability, thaw the vial and initiate the culture as soon as possible upon receipt. If upon arrival, continued storage of the frozen culture is necessary, it should be stored in liquid nitrogen vapor phase and not at -70°C. Storage at -70°C will result in loss of viability.

- Thaw the vial by gentle agitation in a 37°C water bath. To reduce the possibility of contamination, keep the O-ring and cap out of the water. Thawing should be rapid (approximately 2 - 3 minutes).
- Remove the vial from the water bath as soon as the contents are thawed, and decontaminate by dipping in or spraying with 70% ethanol. All of the operations from this point on should be carried out under strict aseptic conditions.
- Transfer the vial contents to a centrifuge tube containing 5.0 mL complete culture medium and spin at approximately 176 x g for 5 minutes. Discard supernatant.
- Resuspend cell pellet with the recommended recovery medium. And dispense into appropriate culture dishes.

- e) Incubate the culture at 37°C in a suitable incubator. A 5% CO<sub>2</sub> in air atmosphere is recommended if using the medium described on this product sheet.

## Cell Freezing

Freezing Medium: 90% FBS+10% DMSO

- Centrifuge at 176 x g for 3 minutes to collect cells.
- Resuspend the cells in pre-cooled freezing medium and adjust the cell density to 5E6 cells/mL.
- Aliquot 1 mL into each vial.
- Place the vial in a controlled-rate freezing container and store at -80°C for at least 1 day, then transfer to liquid nitrogen as soon as possible.

## Cell passage

Growth medium: DMEM+10% FBS+1% P.S+0.75 µg/mL Puromycin

For the first 1 to 2 passages post-resuscitation, use the recovery medium. Once the cells have stabilized, switch to a growth medium.

- Subculturing is necessary when the cell density reaches 80%. It is recommended to perform subculturing at a ratio of 1:3 to 1:4 every 2-3 days. Ensure that the density does not exceed 80%, as overcrowding can lead to reduced viability due to compression.
- Remove and discard culture medium.
- Briefly rinse the cell layer with PBS to remove all traces of serum that contains trypsin inhibitor.
- Add 1.0 mL of 0.25% (w/v) Trypsin-EDTA solution to dish and observe cells under an inverted microscope until cell layer is dispersed (usually within 30 to 60 seconds at 37°C).
- Note: To avoid clumping do not agitate the cells by hitting or shaking the flask while waiting for the cells to detach. Cells that are difficult to detach may be placed at 37°C to facilitate dispersal.
- Add 2.0 mL of growth medium to mix well and aspirate cells by gently pipetting.
- After centrifugation, resuspend the pellet and add appropriate aliquots of the cell suspension to new culture vessels.
- Incubate cultures at 37°C.

**Subcultivation Ratio: A subcultivation ratio of 1:3 - 1:4 is recommended**

**Medium Renewal: Every 2 to 3 days**

## Notes

- Upon initial thawing, a higher number of dead cells is observed, which is a normal phenomenon. Significant improvement is seen after adaptation. Once the cells reach a stable state, the number of dead cells decreases after subculturing and the cell growth rate becomes stable.
- Ensure that the cell density does not exceed 80%, as overcrowding may lead to reduced viability due to compression.

## Sequence

### GUCY2C G7PJX5

MKTLLLDLVLWLLFQPEWLYLTSQVSNCHNGSYEISVLMMDNSAFAEPLNVEDAVNEGLEIVRGRLQN  
 AGLNVTVNASFMYSDDLHNSGDCRSSTCEGLDLLRKISNAKRMGCVLMGSPCTYSTFQMYLDTELSYPMIS  
 AGSFGLSCDYKETLRLMSPARKLTYFLVNFWKTNLDPFKTYSWSTSYVYKNGTESEDCFWYLNALASVS  
 YFSHELKFLVLRQDKEFQDILMDHNRKSNVIVMCGDPEFLYKLGDRVAEDIVIILVDFNDQYFEDNVT  
 APDYMKNVLLVTRSPGNSLLNSSFSRNLSPTRKDFALAYLNGILLFGHMLKTFLNENITTPKFAHAFRNL  
 FEGYDGPVTLDDWGDVDSTMVLLYTSVDTKKYKVLTYDTHVNQTNPVDMSPTFTWKNKLPNDITDRGP  
 QILMIAVFTLTGAVVLLLLVALLMLRKYKDYELRQKKWSHIPPENIFLETNETNHVSLKIDDDKRRDTIQR  
 LRQCKYDKKRVLKDLKHNDGNFTEKQKIELNKLQIDYYNLTKFYGTVKLDTMIFGVIEYCERGLREVLN  
 DTISYPDGTDFMDWEFKISVLYDIAKGMSYLHSSKTEVHGRLKSTNCVVD SRMVVKITDFGCNSILPPKDLW  
 TAPEHLRQANVSQKGDVYSYGIIAQEIILRKETFYTSSCRDRNEKIFRVENSNGMKPFRPDLFLETABEKELEV  
 YLLVKSCWEEDPEKRPDFKKIETTLAKIFGLFHDQKNESYMDTLIRRLQLYSRNLEHLVEERTQLYKAERDRA  
 DRLNFMLLPRLVVKSLKEKGFVEPELYEEVTIYFSDIVGFTTICKYSTPMEVVDMLNDIYKSFHDHVDHVDVY  
 KVETIGDAYMVASGLPKRNGNRHAIDIAKMALEILSFMGTFELEHLPLPIWIRIGVHSGPCAAGVVGIMPR  
 YCLFGDVTNTASRMESTGLPLRIHVSGSTIAILKRTECQFLYEVRGETYKGRGNETTYWLTGMKDQKFNLP  
 TPPTVENQQRLQAEFSDMIANSLQKRQAAGIRSQKPRRVASYKKGTTLEYLQLNTTDKESTYF

## Related Products

CLDN18	
<a href="#">Cynomolgus_CLDN18.2-eGFP CHO-K1 Cell Line</a>	<a href="#">H_CLDN18(isoform2)-eGFP 293 Cell Line</a>
<a href="#">H_CLDN18.1-eGFP HEK-293 Cell Line</a>	<a href="#">H_CLDN18.2 MC38 Cell Line</a>
<a href="#">H_CLDN18.2 MKN45 Cell Line(High Expression)</a>	<a href="#">H_CLDN18.2 MKN45 Cell Line(Low Expression)</a>
<a href="#">H_CLDN18.2 MKN45 Cell Line(Medium Expression)</a>	<a href="#">H_CLDN18.2(isoform2) CHO-K1 Cell Line</a>
<a href="#">H_CLDN18.2-eGFP CT-26 Cell Line</a>	<a href="#">Mouse_CLDN18.2-eGFP CHO-K1 Cell Line</a>
<a href="#">Rat_CLDN18.2-eGFP CHO-K1 Cell Line</a>	<a href="#">Rhesus_CLDN18.2-eGFP CHO-K1 Cell Line</a>
<a href="#">Anti-CLDN18.2 hIgG1 Reference Antibody (IMAB362)</a>	<a href="#">Anti-CLDN18.2 hIgG1 Antibody(LM-102)</a>
<a href="#">Anti-CLDN18.2 hIgG1 Antibody(Zolbetuximab)</a>	
TROP2(TACSTD2)	
<a href="#">Cynomolgus_Trop2 CHO-K1 Cell Line</a>	<a href="#">Cynomolgus_TROP2 HEK-293 Cell Line</a>
<a href="#">H_TROP2 CHO-K1 Cell Line</a>	<a href="#">H_TROP2 CT26 Cell Line</a>
<a href="#">H_TROP2 HEK-293 Cell Line</a>	<a href="#">H_TROP2 LLC1 Cell Line</a>
<a href="#">Anti-H_TROP2 hIgG1 Antibody(Datopotamab)</a>	<a href="#">Anti-TROP2 hIgG1 Antibody(Hu2G10-5)</a>
<a href="#">Anti-Trop2 hIgG1 Reference Antibody (Sacbio)</a>	<a href="#">Anti-Trop2 hIgG1 Reference Antibody(Datbio)</a>
<a href="#">Anti-Trop2-DXD ADC(Dar4)[Datopotamab deruxtecán,Dato-DXD]</a>	<a href="#">Anti-Trop2-SN38 ADC(Dar8)[Sacituzumab govitecan]</a>
<a href="#">Biotinylated Cynomolgus TROP2 Protein; His-Avi Tag</a>	<a href="#">Biotinylated Human TROP2 Protein; His-Avi Tag</a>
<a href="#">Human TROP2 Protein; His Tag</a>	
GUCY2C(GC-C)	
<a href="#">H_GUCY2C CHO-K1 Cell Line</a>	<a href="#">H_GUCY2C HEK-293 Cell Line</a>
<a href="#">Anti-H_GUCY2C hIgG1 Antibody(Indusatumab)</a>	

Human GUCY2C Protein; His Tag	
<b>FGFR2</b>	
H_FGFR2b Reporter 293 Cell Line	Cynomolgus_FGFR2b HEK-293 Cell Line
H_FGFR2b CHO-K1 Cell Line	H_FGFR2b HEK-293 Cell Line
Mouse_FGFR2b HEK-293 Cell Line	
Anti-FGFR2 hIgG1 Antibody(hFR2-14_H12-L1)	Anti-FGFR2 hIgG1 Referece Antibody (Bemabio)
Cynomolgus FGFR2(IIIb) D1-D3 Protein; His Tag	Human FGFR2(IIIb) D1-D3 Protein; hFc Tag
Human FGFR2(IIIb) D2-D3 Protein; hFc Tag	Human FGFR2(IIIb) D2-D3 Protein; His Tag
Human FGFR2(IIIc) D1-D3 Protein; His Tag	Mouse FGFR2(IIIb) D2-D3 Protein; His Tag
<b>CD44</b>	
H_CD44s CHO-K1 Cell Line	H_CD44v6 CHO-K1 Cell Line
H_CD44v6 CHO-K1 Cell Line (Medium Expression)	H_CD44v6 HEK-293 Cell Line
Anti-CD44v6 hIgG1 Antibody(bivatuzumab)	
<b>ADC Related Product</b>	
Anti-DXD Mouse IgG1 Antibody (23E21C5)	Anti-DXD Mouse IgG1 Antibody (4A5A12)
Anti-Dxd Mouse IgG2a Antibody (17D6A4)	Anti-Eribulin Mouse IgG2a Antibody (10F8G4)
Anti-MMAE Mouse IgG1 Antibody (11C10E3)	Anti-MMAE Mouse IgG2a Antibody (17A1K11)
Anti-MMAE Mouse IgG2a Antibody (8F6A3)	Anti-SN38 Mouse IgG1 Antibody(59H11C7)
Mouse anti Human IgG1-DXD(Dar8)	Mouse anti Human IgG1-MMAE(Dar4)
Human IgG1 Isotype-DXD (Dar8)	Human IgG1 Isotype-Eribulin (Dar4)
Human IgG1 Isotype-MMAE (Dar4)	
Recombinant DT3C Protein	

## License Agreement:

**By purchasing and using this cell line product, the user voluntarily agrees to accept and abide by the following policies:**

- This cell line product is restricted to research use only and shall not be used for any commercial purposes.
- This product is strictly prohibited from being used in the diagnosis or treatment of human or animal diseases, and shall not be directly used in experiments involving humans.
- Users and their contractors engaged for their benefit may use this material and its derivatives only within the agreed research scope; modification of the material is not permitted, nor may it be distributed, sold, transferred, or otherwise provided to any other entity (including affiliates).
- If use beyond the above scope is required, prior written permission from Genomeditech (Shanghai) Co.,Ltd. must be obtained. For details, please contact Genomeditech (Shanghai) Co.,Ltd.