Genomeditech (Shanghai) Co., Ltd.
Order: +86 021-50432825/50432826
Email: service@genomeditech.com

Product Sheet

Membrane bound H_KLK3(AA25-261) CHO-K1 Cell Line

Catalog number: GM-C49419

Version 4.0.26.1016

Specifications

Description	Membrane bound H_KLK3(AA25-261) CHO-K1 Cell Line is a clonal stable CHO-K1 cell line that constitutively expresses the Human KLK3 gene, constructed using lentiviral technology.
Quantity	5E6 Cells per vial,1 mL
Product Format	1 vial of frozen cells
Shipping	Shipped on dry ice
Storage Conditions	Liquid nitrogen immediately upon receipt
Target	Human KLK3
Gene ID/Uniprot ID	P07288-1(AA Ile 25 - Pro 261)
Host Cell	CHO-K1
Recovery Medium	F12K+10% FBS+1% P.S
Growth medium	F12K+10% FBS+1% P.S+4 μg/mL Puromycin
Note	None
Freezing Medium	90% FBS+10% DMSO
Growth properties	Adherent
Growth Conditions	37°C, 5% CO_2
Mycoplasma Testing	The cell line has been screened to confirm the absence of Mycoplasma species.
Safety considerations	Biosafety Level 2
Note	It is recommended to expand the cell culture and store a minimum of 10 vials at an early passage for potential future use.

Genomeditech

Materials

Reagent	Manufacturer/Catalogue No.
F12K	BOSTER/PYG0036
Fetal Bovine Serum	ExCell/FSP500
Pen/Strep	Thermo/15140-122
Puromycin	Genomeditech/GM-040401
Anti-KLK3 mIgG2a Antibody(8G8F5)	Genomeditech/GM-88140AB
NovoCyte Flow Cytometer	Agilent/NovoCyte 2060R
Countess 3 Automated Cell Counter	ThermoFisher/A49891

Data

Expression



SampleID	Geometric Mean : R675-H
CHO-K1 anti-KLK3+APC-2nd Ab	695
CHO-K1 Membrane bound H_KLK3(AA25-261) M_IgG+APC-2nd Ab	783
CHO-K1 Membrane bound H_KLK3(AA25-261) anti-KLK3+APC-2nd Ab	120532

Membrane bound H_KLK3(AA25-261) CHO-K1 Cell Line (Cat. GM-C49419) was determined by flow cytometry using Anti-KLK3 mIgG2a Antibody(8G8F5) (Cat. GM-88140AB).

Genomeditech

Cell Recovery

Recovery Medium: F12K+10% FBS+1% P.S

To insure the highest level of viability, thaw the vial and initiate the culture as soon as possible upon receipt. If upon arrival, continued storage of the frozen culture is necessary, it should be stored in liquid nitrogen vapor phase and not at -70°C. Storage at -70°C will result in loss of viability.

a) Thaw the vial by gentle agitation in a 37°C water bath. To reduce the possibility of contamination, keep the O-ring and cap out of the water. Thawing should be rapid (approximately 2 - 3 minutes).
b) Remove the vial from the water bath as soon as the contents are thawed, and decontaminate by dipping in or spraying with 70% ethanol. All of the operations from this point on should be carried out under strict aseptic conditions.
c) Transfer the vial contents to a centrifuge tube containing 5.0 mL complete culture medium and spin at approximately 176 x g for 5 minutes. Discard supernatant.
d) Resuspend cell pellet with the recommended recovery medium. And dispense into appropriate culture dishes.
e) Incubate the culture at 37°C in a suitable incubator. A 5% CO_2 in air atmosphere is recommended if using the medium described on this product sheet.

Cell Freezing

Freezing Medium: 90% FBS+10% DMSO

a) Centrifuge at 176 x g for 3 minutes to collect cells.
b) Resuspend the cells in pre-cooled freezing medium and adjust the cell density to 5E6 cells/mL.
c) Aliquot 1 mL into each vial.
d) Place the vial in a controlled-rate freezing container and store at -80°C for at least 1 day, then transfer to liquid nitrogen as soon as possible.

Cell passage

Growth medium: F12K+10% FBS+1% P.S+4 μg/mL Puromycin

For the first 1 to 2 passages post-resuscitation, use the recovery medium. Once the cells have stabilized, switch to a growth medium.

a) Remove and discard culture medium.
b) Briefly rinse the cell layer with PBS to remove all traces of serum that contains trypsin inhibitor.
c) Add 1.0 mL of 0.25% (w/v) Trypsin-EDTA solution to dish and observe cells under an inverted microscope until cell layer is dispersed (usually within 2 to 3 minutes at 37°C).
d) Note: To avoid clumping do not agitate the cells by hitting or shaking the flask while waiting for the cells to detach. Cells that are difficult to detach may be placed at 37°C to facilitate dispersal.
e) Add 2.0 mL of growth medium to mix well and aspirate cells by gently pipetting.
f) After centrifugation, resuspend the pellet and add appropriate aliquots of the cell suspension to new culture vessels.
g) Incubate cultures at 37°C.

Subcultivation Ratio: A subcultivation ratio of 1:4 - 1:5 is recommended

Medium Renewal: Every 2 to 3 days

Notes

a) After the stabilization of the cell condition, there will be fewer dead cells post-passage, the cell growth rate will tend to stabilize, cell morphology will become uniform, and the cells will appear robust.

Sequence

KLK3 P07288-1(p.25to261)

IVGGWECEKHSQPWQVLVASRGRAVCGGVLVHPQWVLTAAHCIRNKSVILLGRHSLFHPEDTGQVFQVSHSFPH PLYDMSLLKNRFLRPGDDSSHDLMLLRLSEPAELTDAVKVMDLPTQEPALGTTCYASGWGSIEPEEFLTPKKLQC VDLHVISNDVCAQVHPQKVTKFMLCAGRWTGGKSTCSGDSGGPLVCNGVLQGITSWGSEPCALPERPSLYTKV VHYRKWIKDTIVANP

Related Products

FOLH1(PSMA)	
H_FOLH1(PSMA) KO LNCap Cell Line	Cynomolgus_FOLH1(PSMA) CHO-K1 Cell Line
H_FOLH1(PSMA) CHO-K1 Cell Line	H_FOLH1(PSMA) HEK-293 Cell Line
H_FOLH1(PSMA) PC-3 Cell Line	H_FOLH1(PSMA) RM-1 Cell Line
Anti-FOLH1(PSMA) hIgG1 Antibody(Rosopatamab)	Anti-FOLH1(PSMA) hIgG1 Reference Antibody (Rosobio)
Cynomolgus FOLH1(PSMA) Protein; His Tag	
STEAP1	
H_STEAP1 KO LNCaP Cell Line	Cynomolgus_STEAP1 CHO-K1 Cell Line
H_STEAP1 CHO-K1 Cell Line	H_STEAP1 HEK-293 Cell Line
H_STEAP1 RM-1 Cell Line(High Expression)	H_STEAP1 RM-1 Cell Line(Low Expression)
H_STEAP1B2-Flag CHO-K1 Cell Line	Mouse_STEAP1 CHO-K1 Cell Line
Rat_STEAP1 CHO-K1 Cell Line	
Anti-H_STEAP1 hIgG1 Antibody(Vandortuzumab)	Anti-STEAP1 hIgG1 Reference Antibody (Vandbio)
ADC Related Product	
Anti-DXD Mouse IgG1 Antibody (23E21C5)	Anti-DXD Mouse IgG1 Antibody (4A5A12)
Anti-Dxd Mouse IgG2a Antibody (17D6A4)	Anti-Eribulin Mouse IgG2a Antibody (10F8G4)
Anti-MMAE Mouse IgG1 Antibody (11C10E3)	Anti-MMAE Mouse IgG2a Antibody (17A1K11)
Anti-MMAE Mouse IgG2a Antibody (8F6A3)	Anti-SN38 Mouse IgG1 Antibody(59H11C7)
Mouse anti Human IgG1-DXD(Dar8)	Mouse anti Human IgG1-MMAE(Dar4)
Human IgG1 Isotype-DXD (Dar8)	Human IgG1 Isotype-Eribulin (Dar4)
Human IgG1 Isotype-MMAE (Dar4)	
Recombinant DT3C Protein	
AR(NR3C4)	
H_AR Reporter 293 Cell Line	H_AR HEK-293 Cell Line
KLK2 KLK3	
Membrane bound Cynomolgus_KLK2(AA25-270)-Flag CHO-K1 Cell Line	Membrane bound H_KLK2(AA19-261) 22RV1 Cell Line

Membrane bound H_KLK2(AA19-261) CHO-K1 Cell Line	Membrane bound H_KLK2(AA19-261) HEK-293 Cell Line
Membrane bound H_KLK2(AA19-261) LnCap Cell Line	Membrane bound H_KLK2(AA19-261) Luciferase LnCap Cell Line
Membrane bound H_KLK2(AA25-261) CHO-K1 Cell Line	Membrane bound H_KLK2(AA25-261) CT26 Cell Line
Membrane bound H_KLK2(AA25-261) HEK-293 Cell Line	Membrane bound H_KLK2(AA25-261) MC38 Cell Line (Low Expression)
Anti-KLK2 hIgG1 Antibody(Hu11B6)	
ACP3	
Flag-H_ACP3 HCT116 Cell Line	Flag-H_ACP3 HT-1080 Cell Line
H_ACP3 22Rv1 Cell Line	H_ACP3 CHO-K1 Cell Line
H_ACP3 HCT116 Cell Line	H_ACP3 HEK-293 Cell Line
H_ACP3 HT-1080 Cell Line	H_ACP3 PC-3 Cell Line

License Agreement

By purchasing and using this cell line product, the user voluntarily agrees to accept and abide by the following policies:

- This cell line product is restricted to research use only and shall not be used for any commercial purposes.
- This product is strictly prohibited from being used in the diagnosis or treatment of human or animal diseases, and shall not be directly used in experiments involving humans.
- Users and their contractors engaged for their benefit may use this material and its derivatives only within the agreed research scope; modification of the material is not permitted, nor may it be distributed, sold, transferred, or otherwise provided to any other entity (including affiliates).
- If use beyond the above scope is required, prior written permission from Genomeditech (Shanghai) Co.,Ltd. must be obtained. For details, please contact Genomeditech (Shanghai) Co.,Ltd.

Genomeditech