Genomeditech (Shanghai) Co., Ltd.
Order: +86 021-50432825/50432826
Email: service@genomeditech.com

Product Sheet

H_CCR6 Reporter CHO-K1 Cell Line

Catalog number: GM-C48303

Version 4.0.26.1012

CCR6 (C-C chemokine receptor type 6) is a G protein-coupled receptor (GPCR) expressed primarily in immune cells, including immature dendritic cells, memory T cells, and Th17 cells. It is involved in leukocyte chemotaxis, mucosal immunity, and inflammatory responses. CCR6 binds specifically to the chemokine CCL20 (MIP-3α) and is associated with inflammatory and autoimmune diseases such as inflammatory bowel disease, psoriasis, and rheumatoid arthritis, as it recruits pro-inflammatory cells to inflamed tissues. The receptor also plays a role in tumor immunity, making it a potential target for drug development and immunotherapy research.

H_CCR6 Reporter CHO-K1 Cell Line is a clonal stable cell line constitutively expressing CCR6 along with a luciferase reporter gene. When CCL20 binds to CCR6, downstream signaling is activated, leading to luciferase expression. The luciferase activity reflects the activation level of the signaling pathway and can be used to evaluate the in vitro effects of CCR6-related drugs.



Specifications

Quantity	5E6 Cells per vial,1 mL
Product Format	1 vial of frozen cells
Shipping	Shipped on dry ice
Storage Conditions	Liquid nitrogen immediately upon receipt
Recovery Medium	F12K+10% FBS+1% P.S
Growth medium	F12K+10% FBS+1% P.S+4 μg/mL Blasticidin+100 μg/mL Hygromycin+4 μg/mL Puromycin
Note	None
Freezing Medium	90% FBS+10% DMSO
Growth properties	Adherent
Growth Conditions	37°C, 5% CO_2
Mycoplasma Testing	The cell line has been screened to confirm the absence of Mycoplasma species.
Safety considerations	Biosafety Level 2
Note	It is recommended to expand the cell culture and store a minimum of 10 vials at an early passage for potential future use.

Materials

Reagent	Manufacturer/Catalogue No.
F12K	BOSTER/PYG0036
Fetal Bovine Serum	ExCell/FSP500
Pen/Strep	Thermo/15140-122
Blasticidin	Genomeditech/GM-040404
Puromycin	Genomeditech/GM-040401
Hygromycin	Genomeditech/GM-040403
96 well round cell culture plate	PakGent/CL-PT096
96-well White Opaque Plate	Thermo/236108
96-well U-bottom Plate	Saining/1014010
GMOne-Step 2.0 Luciferase Reporter Gene Assay Kit	Genomeditech/GM-040513
Recombinant Human CCL20/MIP-3 alpha Protein	R&D Systems/360-MP-025/CF
Anti-CCR6 hIgG1 Antibody (hAB6)	Genomeditech/GM-89042AB

Genomeditech

Data

Signaling Pathway



Response to Recombinant Human CCL20/MIP-3 alpha Protein. H_CCR6 Reporter CHO-K1 Cell Line (Cat. GM-C48303) was seeded at a density of 1E4 cells per well in a 96-well plate and incubated overnight. The next day, serial dilutions of the Recombinant Human CCL20/MIP-3 alpha Protein (rndsystems/360-MP-025) were added to the pre-seeded cells. The mixture was incubated for an additional 16 hours. The firefly luciferase activity was measured using the Luciferase Reporter Assay Kit (Genomeditech). The maximum induction fold was approximately [21.7]. Data are shown by drug mass concentration.

Expression



	SampleID	Geometric Mean : R675-H
	CHO-K1 Anti-CCR6+APC 2nd Ab	937
	H_CCR6 Reporter CHO-K1 H_IgG+APC 2nd Ab	934
	H_CCR6 Reporter CHO-K1 Anti-CCR6+APC 2nd Ab	316448

H_CCR6 Reporter CHO-K1 Cell Line (Cat. GM-C48303) was determined by flow cytometry using Anti-CCR6 hIgG1 Antibody (hAB6)(Cat. GM-89042AB).

Cell Recovery

Recovery Medium: F12K+10% FBS+1% P.S

To insure the highest level of viability, thaw the vial and initiate the culture as soon as possible upon receipt. If upon arrival, continued storage of the frozen culture is necessary, it should be stored in liquid nitrogen vapor phase and not at -70°C. Storage at -70°C will result in loss of viability.

a) Thaw the vial by gentle agitation in a 37°C water bath. To reduce the possibility of contamination, keep the O-ring and cap out of the water. Thawing should be rapid (approximately 2 - 3 minutes).
b) Remove the vial from the water bath as soon as the contents are thawed, and decontaminate by dipping in or spraying with 70% ethanol. All of the operations from this point on should be carried out under strict aseptic conditions.
c) Transfer the vial contents to a centrifuge tube containing 5.0 mL complete culture medium and spin at approximately 176 x g for 5 minutes. Discard supernatant.
d) Resuspend cell pellet with the recommended recovery medium. And dispense into appropriate culture dishes.
e) Incubate the culture at 37°C in a suitable incubator. A 5% CO_2 in air atmosphere is recommended if using the medium described on this product sheet.

Cell Freezing

Freezing Medium: 90% FBS+10% DMSO

a) Centrifuge at 176 x g for 3 minutes to collect cells.
b) Resuspend the cells in pre-cooled freezing medium and adjust the cell density to 5E6 cells/mL.
c) Aliquot 1 mL into each vial.
d) Place the vial in a controlled-rate freezing container and store at -80°C for at least 1 day, then transfer to liquid nitrogen as soon as possible.

Cell passage

Growth medium: F12K+10% FBS+1% P.S+4 μg/mL Blasticidin+100 μg/mL Hygromycin+4 μg/mL Puromycin

For the first 1 to 2 passages post-resuscitation, use the recovery medium. Once the cells have stabilized, switch to a growth medium.

a) Remove and discard culture medium.
b) Briefly rinse the cell layer with PBS to remove all traces of serum that contains trypsin inhibitor.
c) Add 1.0 mL of 0.25% (w/v) Trypsin-EDTA solution to dish and observe cells under an inverted microscope until cell layer is dispersed (usually within 2 to 3 minutes at 37°C).
d) Note: To avoid clumping do not agitate the cells by hitting or shaking the flask while waiting for the cells to detach. Cells that are difficult to detach may be placed at 37°C to facilitate dispersal.
e) Add 2.0 mL of growth medium to mix well and aspirate cells by gently pipetting.
f) After centrifugation, resuspend the pellet and add appropriate aliquots of the cell suspension to new culture vessels.
g) Incubate cultures at 37°C.

Subcultivation Ratio: A subcultivation ratio of 1:4 - 1:5 is recommended

Medium Renewal: Every 2 to 3 days

Notes

a) After the stabilization of the cell condition, there will be fewer dead cells post-passage, the cell growth rate will tend to stabilize, cell morphology will become uniform, and the cells will appear robust.

Related Products

CCR6	
Cynomolgus_CCR6 CHO-K1 Cell Line	H_CCR6 CHO-K1 Cell Line
H_CCR6 HEK-293 Cell Line(High Expression)	H_CCR6 HEK-293 Cell Line(Low Expression)
Mouse_CCR6 HEK-293 Cell Line	

License Agreement

By purchasing and using this cell line product, the user voluntarily agrees to accept and abide by the following policies:

- This cell line product is restricted to research use only and shall not be used for any commercial purposes.
- This product is strictly prohibited from being used in the diagnosis or treatment of human or animal diseases, and shall not be directly used in experiments involving humans.
- Users and their contractors engaged for their benefit may use this material and its derivatives only within the agreed research scope; modification of the material is not permitted, nor may it be distributed, sold, transferred, or otherwise provided to any other entity (including affiliates).
- If use beyond the above scope is required, prior written permission from Genomeditech (Shanghai) Co.,Ltd. must be obtained. For details, please contact Genomeditech (Shanghai) Co.,Ltd.

Genomeditech